

อภิธานการ

สัญญาเลขที่ R2563C0



สำนักหอสมุด

รายงานวิจัยฉบับสมบูรณ์

การลดรูปสมการเชิงอนุพันธ์สามัญอันดับสองไปสู่สมการเชิงเส้นโดยการ
แปลงเชิงเส้นทั่วไปและการประยุกต์ใช้

ผู้ช่วยศาสตราจารย์ ดร.สุภาพร สุขเสริญ

ภาควิชาคณิตศาสตร์ คณะวิทยาศาสตร์ มหาวิทยาลัยนเรศวร

สำนักหอสมุด มหาวิทยาลัยนเรศวร

วันลงทะเบียน... 28 ก.พ. 2565

เลขทะเบียน... 1049049

เลขเรียกหนังสือ... ว. คค

424

สนับสนุนโดย

๙๙๙๘

๙๕๖๓

งบประมาณรายได้มหาวิทยาลัยนเรศวร

ปีงบประมาณ 2563

สารบัญ

Executive Summary	1
1. ความสำคัญและที่มาของปัญหา.....	1
2. วัตถุประสงค์	2
3. ระเบียบวิธีวิจัย.....	2
4. แผนการดำเนินงานวิจัย	3
5. ตัวชี้วัดเพื่อประเมินผลสำเร็จของโครงการ.....	3
บทคัดย่อ.....	4
Abstract	5
เนื้อหางานวิจัย	6
1. Introduction.....	6
2. Formulation of the linearization theorems.....	7
3. Some applications	13
4. Conclusion	31
References.....	31
ภาคผนวก	33
บทความสำหรับการเผยแพร่	34
ตารางเปรียบเทียบกิจกรรมที่วางแผนไว้ และกิจกรรมที่ดำเนินการมา	43
ผลที่ได้รับตลอดโครงการ	45

Executive Summary

1. ความสำคัญและที่มาของปัญหา

ปัญหาการทำให้เป็นเชิงเส้น (linearization problem) นับเป็นแขนงที่สำคัญแขนงหนึ่งในสาขาของสมการเชิงอนุพันธ์ (differential equation) ในปัจจุบันนักคณิตศาสตร์ได้ศึกษาและวิจัยในแขนงดังกล่าวกันอย่างต่อเนื่อง ในการคิดค้นทฤษฎีเพื่อหาคำตอบความรู้ใหม่ ๆ นั้นนับว่ามีประโยชน์เป็นอย่างมากต่อทางวิชาการ และการพัฒนาประเทศ เป็นที่ยอมรับว่าทฤษฎีและองค์ความรู้ใหม่ ๆ ที่เกิดจากการวิจัยนั้น นอกจากจะมีประโยชน์อย่างมากในการพัฒนาความรู้เชิงวิชาการในสาขาและแขนงต่าง ๆ นั้นแล้ว บางครั้งยังสามารถนำไปประยุกต์ในสาขาอื่น ๆ และเป็นพื้นฐานสำคัญในการพัฒนาทางวิทยาศาสตร์พื้นฐาน (basic science) ซึ่งเป็นการวิจัยพื้นฐาน (basic research) เพื่อสร้างองค์ความรู้ใหม่ อันถือเป็นพื้นฐานในการพัฒนาประเทศชาติต่อไป

ปัญหาการทำให้เป็นเชิงเส้นนับว่าเป็นแขนงหนึ่งที่สามารถประยุกต์ได้อย่างกว้างขวาง โดยเฉพาะอย่างยิ่งต่อการศึกษาเกี่ยวกับการหาผลเฉลยของสมการ ปัญหาที่มีความสำคัญทางกายภาพส่วนใหญ่มักจะอยู่ในรูปแบบของสมการเชิงอนุพันธ์ไม่เชิงเส้น (nonlinear differential equation) ซึ่งตามปกติการแก้ปัญหาที่อยู่ในรูปแบบไม่เชิงเส้นนั้นจะทำได้ยาก และไม่มีวิธีการหาผลเฉลยที่แท้จริง (exact solution) วิธีการเชิงตัวเลข (numerical method) นิยมนำมาแก้ปัญหา แต่ผลเฉลยที่ได้เป็นเพียงผลเฉลยประมาณค่า (approximate solution) อย่างไรก็ตามผลเฉลยที่แท้จริงมีความน่าสนใจกว่าเพราะสามารถนำไปวิเคราะห์คุณสมบัติของสมการที่จะศึกษาได้ หนึ่งในวิธีการที่ใช้หาผลเฉลยที่แท้จริงนี้ คือ การทำสมการที่สนใจศึกษาให้มีความเป็นเชิงเส้น แล้วหาผลเฉลยโดยตรงจากวิธีการพื้นฐาน ซึ่งผลเฉลยที่ได้จากการแก้สมการเชิงเส้น ยังคงเป็นผลเฉลยของสมการที่มีมาแต่เดิมด้วย ซึ่งวิธีการดังกล่าวเราจำเป็นต้องทำการแปลง (transformation) เพื่อแปลงสมการตั้งต้นให้เป็นสมการเชิงเส้น

การแปลงที่น่าสนใจก็มีหลายรูปแบบด้วยกัน ตัวอย่างเช่น ถ้าการแปลงประกอบด้วยอนุพันธ์ เรา จะเรียกการแปลงนี้ว่า การแปลงแบบแทนเจนต์ (tangent transformation) ถ้าการแปลงขึ้นอยู่กับตัวแปรอิสระและตัวแปรตามเท่านั้น เราจะเรียกการแปลงนี้ว่า การแปลงแบบจุด (point transformation) เราจะเรียกการแปลงแบบแทนเจนต์ที่ซึ่งนิยามด้วยการเปลี่ยนตัวแปรอิสระ ตัวแปรตาม และอนุพันธ์อันดับหนึ่งว่า การแปลงแบบคอนแทคท์ (contact transformation) และยังมีอีกชนิดหนึ่งที่มีเซตของการแปลงจะแตกต่างจากการแปลงที่ได้กล่าวมาข้างต้น เนื่องจากประกอบด้วยพจน์ที่ไม่เฉพาะ (nonlocal term)

$$T = \int G(t, x) dt$$

การแปลงชนิดนี้จะเรียกว่า การแปลงแบบซันด์มันทั่วไป (generalized Sundman transformation) ซึ่งในงานวิจัยนี้ การแปลงที่ถูกเลือกมาใช้คือ การแปลงเชิงเส้นทั่วไป (generalized linearizing transformation) ซึ่งเป็นการขยายการแปลงแบบซันด์มันทั่วไป โดยที่ฟังก์ชัน G ที่เลือกใช้ คือ $G(t, x, x')$

เนื่องจากการทำสมการเชิงอนุพันธ์สามัญอันดับสองให้เป็นเชิงเส้นในโดยใช้การแปลงเชิงเส้นทั่วไป

ยังมีผู้ศึกษาไม่ครอบคลุมทุกกรณี ในงานวิจัยของเราจึงมุ่งศึกษาไปยังกรณีที่เหลือ ซึ่งค้นพบว่าเป็นกรณีที่สามารถนำผลไปประยุกต์ใช้แก้ปัญหาภัยกับหลายสมการไม่เชิงเส้นที่มีอยู่จริงในธรรมชาติ

2. วัตถุประสงค์

2.1 หาเงื่อนไขที่จำเป็นและเพียงพอที่ทำให้สมการเชิงอนุพันธ์สามัญอันดับสองใด ๆ ในรูปแบบ

$$x'' = f(t, x, x')$$

ซึ่งสามารถลดรูปไปสู่สมการเชิงเส้นในรูปแบบ

$$X''(T) = 0$$

โดยใช้การแปลงเชิงเส้นทั่วไป ซึ่งอยู่ในรูปแบบ

$$X = F(t, x), \quad dT = [G_1(t, x)x' + G_2(t, x)]dt, \quad G_1 \neq 0$$

2.2 ทหาการแปลงเชิงเส้น

2.3 หาตัวอย่าง และการประยุกต์ใช้ทฤษฎีบท

2.4 สร้างโปรแกรมสำเร็จรูปในการทดสอบความเป็นเชิงเส้น

3. ระเบียบวิธีวิจัย

3.1 ศึกษาโครงสร้างของสมการเชิงอนุพันธ์สามัญอันดับสอง การแปลงในรูปแบบต่างๆ และ ผลงานวิจัยที่เกี่ยวข้องที่มีนักวิจัยทำมาก่อนหน้านี้ ด้วยการสืบค้นข้อมูลในอินเทอร์เน็ต เอกสาร และตำราจากห้องสมุด

3.2 ค้นคว้าหาเอกสาร ตำรา วารสาร และ เอกสารสิ่งพิมพ์ที่เกี่ยวข้องกับงานวิจัยที่กำลังดำเนินการวิจัยอยู่จากแหล่งข้อมูลต่างๆ

3.3 หาเงื่อนไขที่จำเป็นโดยวิธีการ change of derivatives

3.4 หาเงื่อนไขที่เพียงพอโดยใช้ compatibility theory

3.5 ทหาการแปลงเชิงเส้น

3.6 สร้างโปรแกรมสำเร็จรูปในการทดสอบความเป็นเชิงเส้นโดยใช้โปรแกรม Reduce

3.7 หาตัวอย่าง และการนำไปประยุกต์ใช้

3.8 เขียนและพิมพ์ผลงานวิจัยเพื่อส่งตีพิมพ์

3.9 รายงานสรุปผลโครงการ

4. แผนการดำเนินงานวิจัย

กิจกรรม	เดือนที่												
	1	2	3	4	5	6	7	8	9	10	11	12	
1. ศึกษาโครงสร้างของสมการเชิงอนุพันธ์สามัญ อันดับสอง การแปลงในรูปแบบต่างๆ และ ผลงานวิจัยที่เกี่ยวข้องที่มีนักวิจัยทำมาก่อนหน้า นี้	↔												
2. ค้นคว้าหาเอกสาร ตำรา วารสาร และ เอกสารสิ่งพิมพ์ที่เกี่ยวข้องกับงานวิจัยที่กำลัง ดำเนินการวิจัยอยู่จากแหล่งข้อมูลต่างๆ		↔											
3. คิดค้นและวิจัยเพื่อหาเงื่อนไขที่จำเป็นสำหรับ การทำให้เป็นเชิงเส้น			↔										
4. คิดค้นและวิจัยเพื่อหาเงื่อนไขที่เพียงพอ สำหรับการทำให้เป็นเชิงเส้น					↔								
5. คิดค้นและวิจัยเพื่อหาการแปลงเชิงเส้น							↔						
6. สร้างโปรแกรมสำเร็จรูปในการทดสอบความ เป็นเชิงเส้น								↔					
7. คิดค้นและวิจัยเพื่อหาตัวอย่างและการ ประยุกต์ใช้									↔				
8. เขียนและพิมพ์ผลงานวิจัยเพื่อส่งพิจารณา ตีพิมพ์										↔			
9. รายงานสรุปผลโครงการ												↔	

5. ตัวชี้วัดเพื่อการประเมินผลสำเร็จของโครงการ

ตีพิมพ์ในวารสารวิชาการระดับนานาชาติ (ไม่มีค่า Impact Factor)

บทคัดย่อ

ในงานวิจัยนี้เราได้นำเสนอปัญหาการทำให้เป็นเชิงเส้นของสมการเชิงอนุพันธ์สามัญอันดับสองโดยใช้การแปลงเชิงเส้นทั่วไป เราได้พบรูปแบบที่จำเป็นสำหรับการลดรูปสมการเชิงอนุพันธ์สามัญอันดับสองเป็นสมการเชิงเส้นอย่างง่าย นอกจากนี้เรายังได้ค้นพบเงื่อนไขที่เพียงพอสำหรับการทำให้รูปแบบด้านบนแปลงเป็นเชิงเส้นได้ ตลอดจนกระบวนการสำหรับการได้มาของการแปลงเชิงเส้นถูกแสดงให้เห็นในรูปแบบที่ชัดเจน ยิ่งไปกว่านั้นเราประยุกต์ใช้เกณฑ์การทำให้เป็นเชิงเส้นที่ได้มากับปัญหาที่น่าสนใจของสมการเชิงอนุพันธ์สามัญแบบไม่เชิงเส้นและสมการเชิงอนุพันธ์ย่อยแบบไม่เชิงเส้น ตัวอย่างเช่น สมการรุ่มชูชีพ สมการ Painlevé - Gambier XI สมการสำหรับการแกว่งของความถี่ การแกว่งแบบนอนโพลีโนเมียลในหนึ่งมิติ สมการที่สามารถแปลงเป็นเชิงเส้นโดยการแปลงแบบจุดและการแปลงอันดับมัม สมการ Vakhnenko ปรับปรุงทั่วไป


คำสำคัญ : ปัญหาการทำให้เป็นเชิงเส้น การแปลงเชิงเส้นทั่วไป สมการเชิงอนุพันธ์สามัญไม่เชิงเส้นอันดับสอง



Abstract

In this research, we have proposed the linearization problem of second-order ordinary differential equation under the generalized linearizing transformation. We found the necessary form for reducing the second-order ordinary differential equation to simple linear equation. We also obtained sufficient condition for making the above form to be linear. Further, the procedure of linear transformation within the study is demonstrated in the explicit form. Moreover, we apply the obtained linearization criteria to the interesting problems of nonlinear ordinary differential equations and nonlinear partial differential equations, for examples the parachute equation, the Painlevé - Gambier XI equation, the equation for the variable frequency oscillator, the one-dimensional nonpolynomial oscillator, the equation that can be linearizable by point and Sundman transformations, the modified generalized Vakhnenko equation.

Keyword : Linearization problem, generalized linearizing transformation, nonlinear second-order ordinary differential equation



เนื้อหางานวิจัย

1. Introduction

1.1. Introduction to the research problem and its significance

There has been major interest in the nonlinear problems, since most equations are inherently nonlinear in nature. In general the nonlinear problems are very difficult to solve explicitly. It is of interest to provide general criteria for the linearizability of nonlinear ordinary differential equations, as they can then be reduced to easily solvable equations. Therefore, the approach of investigating nonlinear ordinary differential equations via transforming to simpler ordinary differential equations becomes important and has been quite plentiful in analysis of physical problems. This includes the classical linearization problem of finding transformations that linearize a given ordinary differential equation. The linearization problem has been studied in many aspects. A short review can be found in [1, 2]. The tools commonly used for solving the linearization problem are the transformations such as point transformation, contact transformation, reduction of order, differential substitution, generalized Sundman transformation etc. For this research, we employ the extension of the generalized Sundman transformations.

1.2. Historical review

The linearization problem for a second-order ordinary differential equation was investigated with respect to a generalized Sundman transformation

$$X = F(t, x), \quad dT = G(t, x)dt \quad (1.1)$$

by Duarte et al. [3] earlier. They obtained the form of the linearizable equations and the conditions which allow the second-order ordinary differential equation to be transformed to the free particle equation. A characterization of these equations that can be linearized by means of generalized Sundman transformations in terms of first integral and procedure for construction of linearizing transformations has been given by Muriel and Romero [4]. In [5], Mustafa et al. gave a new characterization of linearizable equations in terms of the coefficients of ordinary differential equation and one auxiliary function. In [6], Nakpim and Meleshko pointed out that the solution of the linearization problem for a second-order ordinary differential equation via the generalized Sundman transformation considered earlier by Duarte et al. using the Laguerre form is not complete.

In this work, we expose a more general transformation, i.e. the extension of the generalized Sundman transformation

$$X = F(t, x), \quad dT = G(t, x, x')dt. \quad (1.2)$$

This transformation was studied in [7–9] where they designated the transformation as the *generalized linearizing transformation*. They showed that this transformation can be utilized to linearize a wider class of nonlinear ordinary differential equations and, in particular, certain equations which cannot be linearized by the non-point and invertible point transformations. If the function G in (1.2) is independent of the variable x' then it becomes a non-point transformation (vide equation (1.1)). On the other hand, if G is a differentiable function then it becomes an invertible point transformation. So, (1.2) is

a unified transformation as it includes non-point and invertible point transformations as special cases. An example of an equation which can be linearized by a transformation of the form (1.2) is given in [8].

In [7], the Chandrasekar, Senthilvelan and Lakshmanan applied a particular class of transformations (1.2), where the function $G(t, x, x')$ is linear with respect to x' .

They paid attention to the case where G is a polynomial function in x' and in particular where it is linear in x' with coefficients which are arbitrary functions of t and x . To be specific, they focused here on the case

$$X = F(t, x), \quad dT = (G_1(t, x)x' + G_2(t, x)) dt.$$

Notice that for the case $G_1 = 0$, the generalized linearizing transformation becomes a generalized Sundman transformation, so that they assumed $G_1 \neq 0$.

The authors of [7] obtained that any second-order linearizable ordinary differential equation which can be mapped into the equation $X'' = 0$ via a generalized linearizing transformation has to be of the form

$$x'' + A_3(t, x)x'^3 + A_2(t, x)x'^2 + A_1(t, x)x' + A_0(t, x) = 0, \quad (1.3)$$

and the functions A_i 's ($i = 0, 1, 2, 3$) are connected to the transform functions F and G through the relations

$$\begin{aligned} A_3 &= (G_1 F_{xx} - F_x G_{1x})/M, \\ A_2 &= (G_2 F_{xx} + 2G_1 F_{xt} - F_x G_{2x} - F_t G_{1x} - F_x G_{1t})/M, \\ A_1 &= (2G_2 F_{xt} + G_1 F_{tt} - F_x G_{2t} - F_t G_{2x} - F_t G_{1t})/M, \\ A_0 &= (G_2 F_{tt} - F_t G_{2t})/M \end{aligned} \quad (1.4)$$

with $M = F_x G_2 - F_t G_1 \neq 0$.

They have analyzed a particular case of equation (1.3), namely, $A_3 = 0$ and $A_2 = 0$ in equation (1.4). Complete analysis of the compatibility of arising equations is given for the case $F_x \neq 0$.

In this research, we will analyze, the linearized criteria for a general case of equation (1.3) with the function in the case $F_x = 0$.

2. Formulation of the linearization theorems

2.1. Obtaining necessary condition of linearization

We begin with investigating the necessary conditions for linearization. We consider the second-order ordinary differential equation

$$x'' = F(t, x, x') \quad (2.1)$$

which can be transformed to a simplest linear equation

$$X'' = 0 \quad (2.2)$$

under the generalized linearizing transformation

$$\begin{aligned} X &= F(t, x), \\ dT &= [G_1(t, x)x' + G_2(t, x)] dt, \end{aligned} \quad (2.3)$$

where $G_1 \neq 0$. So that we arrive at the following theorem.

Theorem 2.1. Any second-order ordinary differential equations (2.1) obtained from a linear equation (2.2) by a generalized linearizing transformation (2.3) has to be the form

$$x'' + A_3(t, x)x'^3 + A_2(t, x)x'^2 + A_1(t, x)x' + A_0(t, x) = 0, \quad (2.4)$$

where

$$A_3 = (-F_{xx}G_1 + F_xG_{1x})/(F_tG_1 - F_xG_2), \quad (2.5)$$

$$A_2 = (-2F_{tx}G_1 + F_tG_{1x} - F_{xx}G_2 + F_xG_{1t} + F_xG_{2x})/(F_tG_1 - F_xG_2), \quad (2.6)$$

$$A_1 = (-2F_{tx}G_2 - F_{tt}G_1 + F_tG_{1t} + F_tG_{2x} + F_xG_{2t})/(F_tG_1 - F_xG_2), \quad (2.7)$$

$$A_0 = (-F_{tt}G_2 + F_tG_{2t})/(F_tG_1 - F_xG_2). \quad (2.8)$$

Proof. Applying a generalized linearizing transformation (2.3), one obtains the following transformations

$$\begin{aligned} X'(T) &= \frac{D_t F(t, x)}{D_t \int [G_1(t, x)x' + G_2(t, x)] dt} \\ &= \frac{F_t + x'F_x}{G_1x' + G_2} \\ &= P(t, x, x'), \end{aligned}$$

$$\begin{aligned} X''(T) &= \frac{D_t P}{D_t \int [G_1(t, x)x' + G_2(t, x)] dt} \\ &= \frac{P_t + P_x x' + P_{x'} x''}{G_1 x' + G_2}, \end{aligned}$$

where

$$P_t = \frac{F_{tt}(G_1x' + G_2) - F_t(G_1x'' + G_{1t}x' + G_{2t})}{(G_1x' + G_2)^2},$$

$$P_x = \frac{F_{tx}(G_1x' + G_2) - F_t(G_{1x}x' + G_{2x})}{(G_1x' + G_2)^2},$$

$$P_{x'} = -\frac{F_t G_{1x}''}{(G_1x' + G_2)^2},$$

and $D_t = \frac{\partial}{\partial t} + x' \frac{\partial}{\partial x} + x'' \frac{\partial}{\partial x'} + \dots$ is a total derivative. Substituting the resulting expression in linear equation (2.2), we get the equation (2.4).

2.2. Obtaining sufficient conditions of linearization and linearizing transformation

For obtaining sufficient conditions of linearizability of equation (2.4), one has to solve the compatibility problem of the system of equations (2.5)-(2.8), considering it as overdetermined system of partial differential equations for the functions F, G_1 and G_2 with given coefficients A_i of equation (2.4).

The compatibility analysis depends on the value of F_x . A complete study of all cases is cumbersome. Here a complete solution is given for the case where $F_x = 0$. For convenience of calculations, we set

$$G_3 = \frac{G_2}{G_1}.$$

So that system of equations (2.5)-(2.8) become

$$A_3 = (-F_{xx}G_1 + F_xG_{1x})/(G_1(F_t - F_xG_3)), \quad (2.9)$$

$$A_2 = (-2F_{tx}G_1 + F_tG_{1x} - F_{xx}G_1G_3 + F_xG_{1t} + F_xG_{1x}G_3 + F_xG_{3x}G_1)/(G_1(F_t - F_xG_3)), \quad (2.10)$$

$$A_1 = (-2F_{tx}G_1G_3 - F_{tt}G_1 + F_tG_{1t} + F_tG_{1x}G_3 + F_tG_{3x}G_1 + F_xG_{1t}G_3 + F_xG_{3t}G_1)/(G_1(F_t - F_xG_3)), \quad (2.11)$$

$$A_0 = (-F_{tt}G_1G_3 + F_tG_{1t}G_3 + F_tG_{3t}G_1)/(G_1(F_t - F_xG_3)). \quad (2.12)$$

According to the notation $K = G_1(F_xG_3 - F_t)$, we define the derivative F_t as

$$F_t = (F_xG_1G_3 - K)/G_1. \quad (2.13)$$

Solving equations (2.9)-(2.12) with respect to F_{xx} , K_x , K_t and G_{3t} , one finds

$$F_{xx} = (F_xG_{1x} + A_3K)/G_1, \quad (2.14)$$

$$K_x = (-F_xG_{1t}G_1 + F_xG_{1x}G_1G_3 + F_xG_{3x}G_1^2 + 3G_{1x}K - A_2G_1K + 3A_3G_1G_3K)/(2G_1), \quad (2.15)$$

$$K_t = (-F_xG_{1t}G_1G_3 + F_xG_{1x}G_1G_3^2 + F_xG_{3x}G_1^2G_3 + 4G_{1t}K - G_{1x}G_3K + 2G_{3x}G_1K - 2A_1G_1K + 3A_2G_1G_3K - 3A_3G_1G_3^2K)/(2G_1), \quad (2.16)$$

$$G_{3t} = G_{3x}G_3 + A_0 - A_1G_3 + A_2G_3^2 - A_3G_3^3. \quad (2.17)$$

Comparing the mixed derivative $(K_x)_t = (K_t)_x$, one obtains

$$\begin{aligned} G_{3xx} = & (2A_{0x}F_xG_1^3 - 2A_{1x}F_xG_1^3G_3 + 4A_{1x}G_1^2K - 2A_{2t}G_1^2K \\ & + 2A_{2x}F_xG_1^3G_3^2 - 6A_{2x}G_1^2G_3K + 6A_{3t}G_1^2G_3K - 2A_{3x}F_xG_1^3G_3^3 \\ & + 6A_{3x}G_1^2G_3^2K + 4F_xG_{1tx}G_1^2G_3 - 2F_xG_{1tt}G_1^2 + 3F_xG_{1t}^2G_1 \\ & - 6F_xG_{1t}G_{1x}G_1G_3 + 2F_xG_{1t}G_{3x}G_1^2 - 2F_xG_{1t}A_1G_1^2 \\ & + 4F_xG_{1t}A_2G_1^2G_3 - 6F_xG_{1t}A_3G_1^2G_3^2 - 2F_xG_{1xx}G_1^2G_3^2 \\ & + 3F_xG_{1x}^2G_1G_3^2 - 2F_xG_{1x}G_{3x}G_1^2G_3 + 2F_xG_{1x}A_0G_1^2 \\ & - 2F_xG_{1x}A_2G_1^2G_3^2 + 4F_xG_{1x}A_3G_1^2G_3^3 - F_xG_{3x}^2G_1^3 - 2G_{1tx}G_1K \\ & + 3G_{1t}G_{1x}K - G_{1t}A_2G_1K + 3G_{1t}A_3G_1G_3K + 2G_{1xx}G_1G_3K \\ & - 3G_{1x}^2G_3K + G_{1x}G_{3x}G_1K + G_{1x}A_2G_1G_3K - 3G_{1x}A_3G_1G_3^2K \\ & - 5G_{3x}A_2G_1^2K + 15G_{3x}A_3G_1^2G_3K + 6A_0A_3G_1^2K \\ & - 6A_1A_3G_1^2G_3K + 6A_2A_3G_1^2G_3^2K - 6A_3^2G_1^2G_3^3K)/(4G_1^2K). \end{aligned} \quad (2.18)$$

Comparing the mixed derivatives $(F_{xx})_t = (F_t)_{xx}$ and $(G_{3xx})_t = (G_{3t})_{xx}$, one arrives at the equations

$$\begin{aligned} & -2A_{0x}F_x^2G_1^4 + 2A_{1x}F_x^2G_1^4G_3 - 4A_{1x}F_xG_1^3K + 2A_{2t}F_xG_1^3K \\ & - 2A_{2x}F_x^2G_1^4G_3^2 + 6A_{2x}F_xG_1^3G_3K - 4A_{2x}G_1^2K^2 - 6A_{3t}F_xG_1^3G_3K \\ & + 8A_{3t}G_1^2K^2 + 2A_{3x}F_x^2G_1^4G_3^3 - 6A_{3x}F_xG_1^3G_3^2K + 4A_{3x}G_1^2G_3K^2 \end{aligned}$$

$$\begin{aligned}
& -4F_x^2 G_{1tx} G_1^3 G_3 + 2F_x^2 G_{1tt} G_1^3 - 3F_x^2 G_{1t}^2 G_1^2 + 6F_x^2 G_{1t} G_{1x} G_1^2 G_3 \\
& -2F_x^2 G_{1t} G_{3x} G_1^3 + 2F_x^2 G_{1t} A_1 G_1^3 - 4F_x^2 G_{1t} A_2 G_1^3 G_3 + 6F_x^2 G_{1t} A_3 G_1^3 G_3^2 \\
& + 2F_x^2 G_{1xx} G_1^3 G_3^2 - 3F_x^2 G_{1x}^2 G_1^2 G_3^2 + 2F_x^2 G_{1x} G_{3x} G_1^3 G_3 - 2F_x^2 G_{1x} A_0 G_1^3 \\
& + 2F_x^2 G_{1x} A_2 G_1^3 G_3^2 - 4F_x^2 G_{1x} A_3 G_1^3 G_3^2 + F_x^2 G_{3x}^2 G_1^4 + 6F_x G_{1tx} G_1^2 K \\
& - 9F_x G_{1t} G_{1x} G_1 K + 3F_x G_{1t} A_2 G_1^2 K - 9F_x G_{1t} A_3 G_1^2 G_3 K \\
& - 6F_x G_{1xx} G_1^2 G_3 K + 9F_x G_{1x}^2 G_1 G_3 K - 3F_x G_{1x} G_{3x} G_1^2 K \\
& - 3F_x G_{1x} A_2 G_1^2 G_3 K + 9F_x G_{1x} A_3 G_1^2 G_3^2 K + 3F_x G_{3x} A_2 G_1^3 K \\
& - 9F_x G_{3x} A_3 G_1^3 G_3 K - 6F_x A_0 A_3 G_1^3 K + 6F_x A_1 A_3 G_1^3 G_3 K \\
& - 6F_x A_2 A_3 G_1^3 G_3^2 K + 6F_x A_3^2 G_1^3 G_3^2 K + 4G_{1t} A_3 G_1 K^2 + 4G_{1xx} G_1 K^2 \\
& - 6G_{1x}^2 K^2 - 4G_{1x} A_3 G_1 G_3 K^2 + 8G_{3x} A_3 G_1^2 K^2 - 8A_1 A_3 G_1^2 K^2 \\
& + 2A_2^2 G_1^2 K^2 + 4A_2 A_3 G_1^2 G_3 K^2 - 6A_3^2 G_1^2 G_3 K^2 = 0, \tag{2.19}
\end{aligned}$$

$$\begin{aligned}
& 4A_{0tx} F_x G_1^4 + 4A_{0t} F_x G_{1x} G_1^3 + 12A_{0t} A_3 G_1^3 K - 4A_{0xx} F_x G_1^4 G_3 \\
& - 8A_{0xx} G_1^3 K + 2A_{0x} F_x G_{1t} G_1^3 - 6A_{0x} F_x G_{1x} G_1^3 G_3 - 18A_{0x} F_x G_{3x} G_1^4 \\
& + 8A_{0x} F_x A_1 G_1^4 - 16A_{0x} F_x A_2 G_1^4 G_3 + 24A_{0x} F_x A_3 G_1^4 G_3^2 - 8A_{0x} A_2 G_1^3 K \\
& + 12A_{0x} A_3 G_1^3 G_3 K - 4A_{1tx} F_x G_1^4 G_3 + 8A_{1tx} G_1^3 K - 4A_{1t} F_x G_{1t} G_1^3 \\
& - 12A_{1t} A_3 G_1^3 G_3 K + 4A_{1xx} F_x G_1^4 G_3^2 + 2A_{1x} F_x G_{1t} G_1^3 G_3 + 2A_{1x} F_x G_{1x} G_1^3 G_3^2 \\
& + 18A_{1x} F_x G_{3x} G_1^4 G_3 - 4A_{1x} F_x A_0 G_1^4 - 4A_{1x} F_x A_1 G_1^4 G_3 + 12A_{1x} F_x A_2 G_1^4 G_3^2 \\
& - 20A_{1x} F_x A_3 G_1^4 G_3^2 - 8A_{1x} G_{3x} G_1^3 K + 8A_{1x} A_1 G_1^3 K - 8A_{1x} A_2 G_1^3 G_3 K \\
& + 12A_{1x} A_3 G_1^3 G_3^2 K + 4A_{2tx} F_x G_1^4 G_3^2 - 8A_{2tx} G_1^3 G_3 K - 4A_{2tt} G_1^3 K \\
& + 8A_{2t} F_x G_{1t} G_1^3 G_3 - 4A_{2t} F_x G_{1x} G_1^3 G_3^2 - 2A_{2t} G_{1t} G_1^2 K + 2A_{2t} G_{1x} G_1^2 G_3 K \\
& + 2A_{2t} G_{3x} G_1^3 K - 4A_{2t} A_1 G_1^3 K + 8A_{2t} A_2 G_1^3 G_3 K - 4A_{2xx} F_x G_1^4 G_3^2 \\
& + 4A_{2xx} G_1^3 G_3^2 K - 6A_{2x} F_x G_{1t} G_1^3 G_3^2 + 2A_{2x} F_x G_{1x} G_1^3 G_3^2 - 18A_{2x} F_x G_{3x} G_1^4 G_3^2 \\
& + 8A_{2x} F_x A_0 G_1^4 G_3 - 8A_{2x} F_x A_2 G_1^4 G_3^2 + 16A_{2x} F_x A_3 G_1^4 G_3^2 + 2A_{2x} G_{1t} G_1^2 G_3 K \\
& - 2A_{2x} G_{1x} G_1^2 G_3^2 K + 14A_{2x} G_{3x} G_1^3 G_3 K - 12A_{2x} A_0 G_1^3 K + 4A_{2x} A_2 G_1^3 G_3^2 K \\
& + 8A_{3t} F_x G_{1x} G_1^3 G_3^2 + 6A_{3t} G_{1t} G_1^2 G_3 K - 6A_{3t} G_{1x} G_1^2 G_3^2 K - 6A_{3t} G_{3x} G_1^3 G_3 K \\
& + 24A_{3t} A_0 G_1^3 K - 12A_{3t} A_1 G_1^3 G_3 K + 4A_{3xx} F_x G_1^4 G_3^2 - 4A_{3xx} G_1^3 G_3^2 K \\
& + 10A_{3x} F_x G_{1t} G_1^3 G_3^2 - 6A_{3x} F_x G_{1x} G_1^3 G_3^2 + 18A_{3x} F_x G_{3x} G_1^4 G_3^2 \\
& - 12A_{3x} F_x A_0 G_1^4 G_3^2 + 4A_{3x} F_x A_1 G_1^4 G_3^2 + 4A_{3x} F_x A_2 G_1^4 G_3^2 \\
& - 12A_{3x} F_x A_3 G_1^4 G_3^2 - 6A_{3x} G_{1t} G_1^2 G_3^2 K + 6A_{3x} G_{1x} G_1^2 G_3^2 K \\
& - 18A_{3x} G_{3x} G_1^3 G_3^2 K + 12A_{3x} A_0 G_1^3 G_3 K - 4A_{3x} A_2 G_1^3 G_3^2 K \\
& + 12A_{3x} A_3 G_1^3 G_3^2 K - 12F_x G_{1txx} G_1^3 G_3^2 - 36F_x G_{1tx} G_{1t} G_1^2 G_3 \\
& + 36F_x G_{1tx} G_{1x} G_1^2 G_3^2 - 36F_x G_{1tx} G_{3x} G_1^3 G_3 + 12F_x G_{1tx} A_0 G_1^3 \\
& + 12F_x G_{1tx} A_1 G_1^3 G_3 - 36F_x G_{1tx} A_2 G_1^3 G_3^2 + 60F_x G_{1tx} A_3 G_1^3 G_3^2 - 4F_x G_{1ttt} G_1^3
\end{aligned}$$

$$\begin{aligned}
& + 12F_x G_{1ttx} G_1^3 G_3 + 18F_x G_{1tt} G_{1t} G_1^2 - 18F_x G_{1tt} G_{1x} G_1^2 G_3 + 18F_x G_{1tt} G_{3x} G_1^3 \\
& - 12F_x G_{1tt} A_1 G_1^3 + 24F_x G_{1tt} A_2 G_1^3 G_3 - 36F_x G_{1tt} A_3 G_1^3 G_3^2 - 15F_x G_{1t}^3 G_1 \\
& + 45F_x G_{1t}^2 G_{1x} G_1 G_3 - 27F_x G_{1t}^2 G_{3x} G_1^2 + 18F_x G_{1t}^2 A_1 G_1^2 - 36F_x G_{1t}^2 A_2 G_1^2 G_3 \\
& + 54F_x G_{1t}^2 A_3 G_1^2 G_3^2 + 18F_x G_{1t} G_{1xx} G_1^2 G_3^2 - 45F_x G_{1t} G_{1x}^2 G_1 G_3^2 \\
& + 54F_x G_{1t} G_{1x} G_{3x} G_1^2 G_3 - 18F_x G_{1t} G_{1x} A_0 G_1^2 - 18F_x G_{1t} G_{1x} A_1 G_1^2 G_3 \\
& + 54F_x G_{1t} G_{1x} A_2 G_1^2 G_3^2 - 90F_x G_{1t} G_{1x} A_3 G_1^2 G_3^3 - 9F_x G_{1t} G_{3x}^2 G_1^3 \\
& + 18F_x G_{1t} G_{3x} A_1 G_1^3 - 36F_x G_{1t} G_{3x} A_2 G_1^3 G_3 + 54F_x G_{1t} G_{3x} A_3 G_1^3 G_3^2 \\
& + 8F_x G_{1t} A_0 A_2 G_1^3 - 24F_x G_{1t} A_0 A_3 G_1^3 G_3 - 8F_x G_{1t} A_1^2 G_1^3 \\
& + 24F_x G_{1t} A_1 A_2 G_1^3 G_3 - 24F_x G_{1t} A_1 A_3 G_1^3 G_3^2 - 24F_x G_{1t} A_2^2 G_1^3 G_3^2 \\
& + 64F_x G_{1t} A_2 A_3 G_1^3 G_3^3 - 48F_x G_{1t} A_3^2 G_1^3 G_3^4 + 4F_x G_{1xxx} G_1^3 G_3^3 \\
& - 18F_x G_{1xx} G_{1x} G_1^3 G_3^3 + 18F_x G_{1xx} G_{3x} G_1^3 G_3^2 - 12F_x G_{1xx} A_0 G_1^3 G_3 \\
& + 12F_x G_{1xx} A_2 G_1^3 G_3^3 - 24F_x G_{1xx} A_3 G_1^3 G_3^4 + 15F_x G_{1x}^3 G_1 G_3^3 \\
& - 27F_x G_{1x}^2 G_{3x} G_1^2 G_3^2 + 18F_x G_{1x}^2 A_0 G_1^2 G_3 - 18F_x G_{1x}^2 A_2 G_1^2 G_3^3 \\
& + 36F_x G_{1x}^2 A_3 G_1^2 G_3^4 + 9F_x G_{1x} G_{3x}^2 G_1^3 G_3 - 18F_x G_{1x} G_{3x} A_0 G_1^3 \\
& + 18F_x G_{1x} G_{3x} A_2 G_1^3 G_3^2 - 36F_x G_{1x} G_{3x} A_3 G_1^3 G_3^3 + 8F_x G_{1x} A_0 A_1 G_1^3 \\
& - 24F_x G_{1x} A_0 A_2 G_1^3 G_3 + 48F_x G_{1x} A_0 A_3 G_1^3 G_3^2 - 8F_x G_{1x} A_1 A_3 G_1^3 G_3^3 \\
& + 8F_x G_{1x} A_2^2 G_1^3 G_3^3 - 24F_x G_{1x} A_2 A_3 G_1^3 G_3^4 + 24F_x G_{1x} A_3^2 G_1^3 G_3^5 + 3F_x G_{3x}^3 G_1^4 \\
& + 8G_{1txx} G_1^2 G_3 K + 10G_{1tx} G_{1t} G_1 K - 26G_{1tx} G_{1x} G_1 G_3 K + 14G_{1tx} G_{3x} G_1^2 K \\
& - 4G_{1tx} A_1 G_1^2 K + 16G_{1tx} A_2 G_1^2 G_3 K - 36G_{1tx} A_3 G_1^2 G_3^2 K - 4G_{1tx} G_1^2 K \\
& + 8G_{1tt} G_{1x} G_1 K - 4G_{1tt} A_2 G_1^2 K + 12G_{1tt} A_3 G_1^2 G_3 K - 15G_{1t}^2 G_{1x} K \\
& + 5G_{1t}^2 A_2 G_1 K - 15G_{1t}^2 A_3 G_1 G_3 K - 10G_{1t} G_{1xx} G_1 G_3 K + 30G_{1t} G_{1x}^2 G_3 K \\
& - 22G_{1t} G_{1x} G_{3x} G_1 K + 8G_{1t} G_{1x} A_1 G_1 K - 26G_{1t} G_{1x} A_2 G_1 G_3 K \\
& + 54G_{1t} G_{1x} A_3 G_1 G_3^2 K + 8G_{1t} G_{3x} A_2 G_1^2 K - 24G_{1t} G_{3x} A_3 G_1^2 G_3 K \\
& + 6G_{1t} A_0 A_3 G_1^2 K - 4G_{1t} A_1 A_2 G_1^2 K + 6G_{1t} A_1 A_3 G_1^2 G_3 K + 8G_{1t} A_2^2 G_1^2 G_3 K \\
& - 30G_{1t} A_2 A_3 G_1^2 G_3^2 K + 30G_{1t} A_3^2 G_1^2 G_3^3 K - 4G_{1xxx} G_1^2 G_3^2 K \\
& + 18G_{1xx} G_{1x} G_1 G_3^2 K - 14G_{1xx} G_{3x} G_1^2 G_3 K + 4G_{1xx} A_0 G_1^2 K \\
& - 8G_{1xx} A_2 G_1^2 G_3^2 K + 20G_{1xx} A_3 G_1^2 G_3^3 K - 15G_{1x}^3 G_3^2 K + 22G_{1x}^2 G_{3x} G_1 G_3 K \\
& - 8G_{1x}^2 A_0 G_1 K + 13G_{1x}^2 A_2 G_1 G_3^2 K - 31G_{1x}^2 A_3 G_1 G_3^3 K - 3G_{1x} G_{3x}^2 G_1^2 K \\
& - 8G_{1x} G_{3x} A_2 G_1^2 G_3 K + 24G_{1x} G_{3x} A_3 G_1^2 G_3^2 K + 4G_{1x} A_0 A_2 G_1^2 K \\
& - 18G_{1x} A_0 A_3 G_1^2 G_3 K + 6G_{1x} A_1 A_3 G_1^2 G_3^2 K - 4G_{1x} A_2^2 G_1^2 G_3^2 K \\
& + 14G_{1x} A_2 A_3 G_1^2 G_3^3 K - 18G_{1x} A_3^2 G_1^2 G_3^4 K + 3G_{3x}^2 A_2 G_1^3 K \\
& - 9G_{3x}^2 A_3 G_1^3 G_3 K - 6G_{3x} A_0 A_3 G_1^3 K + 6G_{3x} A_1 A_3 G_1^3 G_3 K \\
& - 6G_{3x} A_2 A_3 G_1^3 G_3^2 K + 6G_{3x} A_3^2 G_1^3 G_3^3 K = 0.
\end{aligned} \tag{2.20}$$

Case $F_x = 0$

Since $F_x = 0$, then substituting it into F_{xx} in equation (2.14), one gets the condition

$$A_3 = 0. \quad (2.21)$$

Comparing the mixed derivative $(F_t)_x = (F_x)_t$, one obtains the derivative

$$G_{1x} = A_2G_1 - 3A_3G_1G_3 \quad (2.22)$$

and this satisfies equation (2.19). Setting

$$\begin{aligned} \lambda_1 &= -A_{1x} + 2A_{2t}, \\ \lambda_2 &= -A_{0xx} - A_{0x}A_2 + A_{2tt} + A_{2t}A_1 - A_{2x}A_0 - \lambda_{1t} - A_1\lambda_1 \end{aligned}$$

then, equation (2.20) becomes

$$G_{3x}\lambda_1 + G_3A_2\lambda_1 + \lambda_2 = 0. \quad (2.23)$$

The compatibility analysis depends on the value of λ_1 . A complete study of all cases is given here.

2.2.1. Case $\lambda_1 = 0$

From equation (2.23), one finds

$$\lambda_2 = 0. \quad (2.24)$$

2.2.2. Case $\lambda_1 \neq 0$

Equation (2.23) provides the derivative

$$G_{3x} = -(G_3A_2\lambda_1 + \lambda_2)/\lambda_1. \quad (2.25)$$

Substituting G_{3x} into G_{3xx} in equation (2.18), one arrives at the condition

$$\lambda_{2x} = (-A_{2t}\lambda_1^2 + \lambda_{1x}\lambda_2 + \lambda_1^3)/\lambda_1. \quad (2.26)$$

Comparing the mixed derivatives $(G_{3x})_t = (G_{3t})_x$, one gets the condition

$$\lambda_{2t} = -(A_{0x}\lambda_1^2 - \lambda_{1t}\lambda_2 + A_0A_2\lambda_1^2 + A_1\lambda_1\lambda_2 + \lambda_2^2)/\lambda_1. \quad (2.27)$$

Combining all derived results in the case $F_x = 0$ the following theorems are proven.

Theorem 2.2. *Sufficient conditions for equation (2.4) to be equivalent to a linear equation (2.2) via generalized linearizing transformation (2.3) with the function $F = F(t)$ is the equation (2.21) and the additional conditions are as follows.*

- (a) *If $\lambda_1 = 0$, then the condition is equation (2.24).*
- (b) *If $\lambda_1 \neq 0$, then the conditions are equations (2.26) and (2.27).*

Corollary 2.3. *Provided that the sufficient conditions in Theorem 2.2 are satisfied, the transformation (2.3) with the function $F = F(t)$ mapping equation (2.4) to a linear equation (2.2) is obtained by solving the compatible system of equations :*

- (a) *(2.13), (2.15), (2.16), (2.17), (2.18), and (2.22).*
- (b) *(2.13), (2.15), (2.16), (2.17), (2.22), and (2.25).*

3. Some applications

3.1. Parachute equation

The idea of this application is based on a model for movement of a parachutist during the air using Newton's II law is $\sum F = ma$. The motion of skydiver when the coefficient of air resistance change between free-fall and the final steady state descent with the parachute fully deployed. Consider the parachute equation [4], in the form

$$x'' - kx'^2 + g = 0, \quad (3.1)$$

with initial conditions $x(0) = 0$ and $x'(0) = 0$.

Here $k = \frac{\pi \rho C_d D^2}{8m}$, where

- m is the mass of the body and parachute,
- ρ is the density of the fluid in which the body moves,
- C_d is the drag coefficient for the parachute (1.5 for parabolic profile and 0.75 for flat),
- D is the effective diameter of the parachute.

It is an equation of the form (2.4) in Theorem 3.2 with the coefficients

$$A_3 = 0, A_2 = k, A_1 = 0, A_0 = g, \lambda_1 = 0, \lambda_2 = 0.$$

One can check that these coefficients obey the conditions in Theorem 2.2. case (a). Thus, equation (3.1) is linearizable via a generalized linearizing transformation. For finding the functions F , G_1 and G_2 we have to solve equations in Corollary 2.3 case (a), which become

$$F_t = -\frac{K}{G_1}, \quad (3.2)$$

$$K_x = kK, \quad (3.3)$$

$$K_t = \frac{K(2G_{1t} + G_1 G_3 k)}{G_1}, \quad (3.4)$$

$$G_{3t} = g + G_3^2 k, \quad (3.5)$$

$$G_{3xx} = 0, \quad (3.6)$$

$$G_{1x} = G_1 k. \quad (3.7)$$

From equation (3.7), we have

$$\begin{aligned} \frac{G_{1x}}{G_1} &= k \\ \int \frac{1}{G_1} dG_1 &= \int k dx \\ \ln G_1 &= kx + C(t) \\ G_1 &= C_1(t) e^{kx}. \end{aligned}$$

Choosing $C_1(t) = 1$, one obtains

$$G_1 = e^{kx}.$$

From equations (3.5) and (3.6), there exists $G_3 = \sqrt{\frac{g}{k}}i$ such that these equations are hold. Then, let

$$G_3 = \sqrt{\frac{g}{k}}i$$

so,

$$G_2 = \sqrt{\frac{g}{k}}ie^{kx}.$$

From equation (3.3), we have

$$\begin{aligned}\frac{K_x}{K} &= k \\ \int \frac{1}{K} dK &= \int k dx \\ \ln K &= kx + C(t) \\ K &= C_2(t)e^{kx}.\end{aligned}$$

From equation (3.4), we have

$$\begin{aligned}\frac{K_t}{K} &= \sqrt{kgi} \\ \int \frac{1}{K} dK &= \int \sqrt{kgit} dt \\ \ln K &= \sqrt{kgit} + C(x) \\ K &= C_3(x)e^{\sqrt{kgit}}.\end{aligned}$$

Choosing $C_2(t) = e^{\sqrt{kgit}}$ and $C_3(x) = e^{kx}$, then

$$K = e^{kx + \sqrt{kgit}}.$$

Thus, equation (3.2) becomes

$$\begin{aligned}F_t &= -e^{\sqrt{kgit}} \\ F &= \frac{i}{\sqrt{kg}}e^{\sqrt{kgit}} + C_4(x).\end{aligned}$$

Choosing $C_4(x) = 0$, one gets

$$F = \frac{i}{\sqrt{kg}}e^{\sqrt{kgit}}.$$

So that, one obtains the linearizing transformation

$$X = \frac{i}{\sqrt{kg}}e^{\sqrt{kgit}}, \quad dT = (e^{kx}x' + \sqrt{\frac{g}{k}}ie^{kx})dt. \quad (3.8)$$

Hence, equation (3.1) is mapped by the transformation (3.8) into the linear equation

$$X'' = 0. \quad (3.9)$$

The general solution of equation (3.9) is

$$X = c_1 T + c_2, \quad (3.10)$$

where c_1 and c_2 are arbitrary constants. Applying the generalized linearizing transformation (3.8) to equation (3.10), we obtain that the general solution of equation (3.1) is

$$\frac{i}{\sqrt{kg}} e^{\sqrt{kg}it} = c_1 \phi(t) + c_2,$$

where the function $T = \phi(t)$ is a solution of the equation

$$\frac{dT}{dt} = (x' + \sqrt{\frac{g}{k}} i) e^{kx}.$$

3.2. Painlevé - Gambier XI equation

In [4], Koudahoun, Akande, Adjai, Kpomahou and Monsia considered the Painlevé - Gambier XI equation

$$x'' + \frac{x'^2}{x} = 0. \quad (3.11)$$

They introduced a generalized singular differential equation of quadratic Liénard type for study of exact classical and quantum mechanical solutions.

By using our obtained theorems, we get the results as follow. Equation (3.11) is an equation of the form (2.4) in Theorem 2.1 with the coefficients

$$A_3 = 0, A_2 = \frac{1}{x}, A_1 = 0, A_0 = 0, \lambda_1 = 0, \lambda_2 = 0.$$

One can check that these coefficients obey the conditions in Theorem 2.2. case (a). Thus, equation (3.11) is linearizable via a generalized linearizing transformation. For finding the functions F , G_1 and G_2 we have to solve equations in Corollary 2.3 case (a), which become

$$F_t = -\frac{K}{G_1}, \quad (3.12)$$

$$K_x = \frac{K}{x}, \quad (3.13)$$

$$K_t = \frac{K(2G_{1t}x + G_1G_3)}{G_1x}, \quad (3.14)$$

$$G_{3t} = \frac{G_3^2}{x}, \quad (3.15)$$

$$G_{3xx} = \frac{G_3}{x^2}, \quad (3.16)$$

$$G_{1x} = \frac{G_1}{x}. \quad (3.17)$$

Consider equation (3.17), we have

$$\begin{aligned} \frac{G_{1x}}{G_1} &= \frac{1}{x} \\ \int \frac{1}{G_1} dG_1 &= \int \frac{1}{x} dx \\ \ln G_1 &= \ln x + \ln C(t) \\ G_1 &= C_1(t)x. \end{aligned}$$

Choosing $C_1(t) = 1$, one obtains

$$G_1 = x.$$

From equations (3.15) and (3.16), there exists $G_3 = 0$ such that these equations are hold. Then, let

$$G_3 = 0$$

so,

$$G_2 = 0.$$

Consider equation (3.13),

$$\begin{aligned} \frac{K_x}{K} &= \frac{1}{x} \\ \int \frac{1}{K} dK &= \int \frac{1}{x} dx \\ \ln K &= \ln x + \ln C(t) \\ K &= C_2(t)x. \end{aligned}$$

Consider equation (3.14),

$$\begin{aligned} K_t &= 0 \\ K &= C_3(x). \end{aligned}$$

Choosing $C_2(t) = 1$ and $C_3(x) = x$, then

$$K = x.$$

Thus, equation (3.12) becomes

$$\begin{aligned} F_t &= -1 \\ F &= -t + C_4(x). \end{aligned}$$

Choosing $C_4(x) = 0$, so

$$F = -t.$$

So that, one obtains the linearizing transformation

$$X = -t, dT = xx'dt. \quad (3.18)$$

Hence, equation (3.11) is mapped by the transformation (3.18) into the linear equation

$$X'' = 0. \quad (3.19)$$

The general solution of equation (3.19) is

$$X = c_1T + c_2, \quad (3.20)$$

where c_1 and c_2 , are arbitrary constants. Applying the generalized linearizing transformation (3.18) to equation (3.20), we obtain that the general solution of equation (3.11) is

$$-t = c_1\phi(t) + c_2,$$

where the function $T = \phi(t)$ is a solution of the equation

$$\frac{dT}{dt} = x x'.$$

3.3. Equation for the variable frequency oscillator

In 2013, Mastafa, Al-Dueik and Mara'beh [4] considered the ordinary differential for the variable frequency oscillator

$$x'' + x x'^2 = 0. \quad (3.21)$$

They showed that this equation can be linearizable by generalized Sundman transformation

$$X = F(t, x), dT = G(t, x)dt, F_x \neq 0.$$

By using their method, the solution of equation (3.21) is

$$\operatorname{erfi}\left(\frac{x}{\sqrt{2}}\right) = C_1t + C_2,$$

where $\operatorname{erfi}(x) = \frac{2}{\sqrt{\pi}} \int_0^x e^{v^2} dv$ is an imaginary error function and C_1, C_2 are arbitrary constants.

By using our obtained theorems, we get the results as follow. Equation (3.21) is an equation of the form (2.4) in Theorem 2.1 with the coefficients

$$A_3 = 0, A_2 = x, A_1 = 0, A_0 = 0, \lambda_1 = 0, \lambda_2 = 0.$$

One can check that these coefficients obey the conditions in Theorem 2.2. case (a). Thus, equation (3.21) is linearizable via a generalized linearizing transformation. For finding the functions F , G_1 and G_2 we have to solve equations in Corollary 2.3 case (a), which become

$$F_t = -\frac{K}{G_1}, \quad (3.22)$$

$$K_x = xK, \quad (3.23)$$

$$K_t = \frac{K(2G_{1t} + G_1 G_3 x)}{G_1}, \quad (3.24)$$

$$G_{3t} = G_3^2 x, \quad (3.25)$$

$$G_{3xx} = -G_3, \quad (3.26)$$

$$G_{1x} = G_1 x. \quad (3.27)$$

Consider equation (3.27), we have

$$\begin{aligned} \frac{G_{1x}}{G_1} &= x \\ \int \frac{1}{G_1} dG_1 &= \int x dx \\ \ln G_1 &= \frac{x^2}{2} + C(t) \\ G_1 &= C_1(t) e^{\frac{x^2}{2}}. \end{aligned}$$

Choosing $C_1(t) = 1$, one obtains

$$G_1 = e^{\frac{x^2}{2}}.$$

From equations (3.25) and (3.26), there exists $G_3 = 0$ such that these equations are hold. Then, let

$$G_3 = 0$$

so,

$$G_2 = 0.$$

Consider equation (3.23),

$$\begin{aligned} \frac{K_x}{K} &= x \\ \int \frac{1}{K} dK &= \int x dx \\ \ln K &= \frac{x^2}{2} + C(t) \\ K &= C_2(t) e^{\frac{x^2}{2}}. \end{aligned}$$

Consider equation (3.24),

$$\begin{aligned} K_t &= 0 \\ K &= C_3(x). \end{aligned}$$

Choosing $C_2(t) = 1$ and $C_3(x) = e^{\frac{x^2}{2}}$, then

$$K = e^{\frac{x^2}{2}}.$$

Thus, equation (3.22) becomes

$$F_t = -1$$

$$F = -t + C_4(x).$$

Choosing $C_4(x) = 0$, so

$$F = -t.$$

So that, one obtains the linearizing transformation

$$X = -t, \quad dT = e^{\frac{x^2}{2}} x' dt. \quad (3.28)$$

Hence, equation (3.21) is mapped by the transformation (3.28) into the linear equation

$$X'' = 0. \quad (3.29)$$

The general solution of equation (3.29) is

$$X = c_1 T + c_2, \quad (3.30)$$

where c_1 and c_2 , are arbitrary constants. Applying the generalized linearizing transformation (3.28) to equation (3.30), we obtain that the general solution of equation (3.21) is

$$-t = c_1 \phi(t) + c_2,$$

where the function $T = \phi(t)$ is a solution of the equation

$$\frac{dT}{dt} = e^{\frac{x^2}{2}} x'.$$

3.4. The one-dimensional non-polynomial oscillator

In the note [4], Mathew and Lakshmanan presented a remarkable nonlinear system that all its bounded periodic motions are simple harmonic. The system is a particle obeying the highly nonlinear equation of motion

$$(1 + \lambda x^2)x'' + (\alpha - \lambda x'^2)x = 0, \quad (3.31)$$

where λ and α are arbitrary parameters.

It is an equation of the form (2.4) in Theorem 2.1 with the coefficients

$$A_3 = 0, A_2 = -\frac{\lambda x}{(\lambda x^2 + 1)}, A_1 = 0, A_0 = \frac{\alpha x}{(\lambda x^2 + 1)},$$

$$\lambda_1 = 0, \lambda_2 = \alpha \lambda x (-\lambda x^2 + 2).$$

One can check that the condition (2.21) in Theorem 2.2. case (a) are satisfied. Now, the condition (2.24) is satisfied when the following condition holds, that is,

$$\alpha \lambda x (-\lambda x^2 + 2) = 0.$$

Two cases arise, that are $\alpha = 0$ and $\lambda = \frac{2}{x^2}$. (Note that for $\lambda = 0$ equation (3.31) is linear equation.)

Here we consider only case $\alpha = 0$. In this case, the equation (3.31) takes the form

$$(1 + \lambda x^2)x'' - \lambda x x'^2 = 0. \quad (3.32)$$

The linearizing transformation is found by solving equations in Corollary 2.3 case (a), which become

$$F_t = -\frac{K}{G_1}, \quad (3.33)$$

$$K_x = -\frac{\lambda x K}{(1 + \lambda x^2)}, \quad (3.34)$$

$$K_t = \frac{K(2G_{1t}\lambda x^2 + 2G_{1t} - \lambda x G_1 G_3)}{G_1(1 + \lambda x^2)}, \quad (3.35)$$

$$G_{3t} = -\frac{\lambda x^2 G_3^2}{(1 + \lambda x^2)}, \quad (3.36)$$

$$G_{3xx} = \frac{\lambda G_3(-\lambda x^2 + 1)}{(1 + \lambda x^2)^2}, \quad (3.37)$$

$$G_{1x} = -\frac{\lambda x G_1}{(1 + \lambda x^2)}. \quad (3.38)$$

Consider equation (3.38), we have

$$\begin{aligned} \frac{G_{1x}}{G_1} &= -\frac{\lambda x}{(1 + \lambda x^2)} \\ \int \frac{1}{G_1} dG_1 &= -\int \frac{\lambda x}{(1 + \lambda x^2)} dx \\ \ln G_1 &= -\frac{1}{2} \ln(1 + \lambda x^2) + \ln C(t) \\ G_1 &= \frac{C_1(t)}{(1 + \lambda x^2)^{\frac{1}{2}}}. \end{aligned}$$

Choosing $C_1(t) = 1$, we obtain

$$G_1 = \frac{1}{(1 + \lambda x^2)^{\frac{1}{2}}}.$$

From equations (3.36) and (3.37), there exists $G_3 = 0$ such that these equations are hold. Then, let

$$G_3 = 0$$

so,

$$G_2 = 0.$$

From equation (3.34), we have

$$\begin{aligned} \frac{K_x}{K} &= -\frac{\lambda x}{(1 + \lambda x^2)} \\ \int \frac{1}{K} dK &= -\int \frac{\lambda x}{(1 + \lambda x^2)} dx \end{aligned}$$

$$\ln K = -\frac{1}{2} \ln(1 + \lambda x^2) + \ln C(t)$$

$$K = \frac{C_2(t)}{(1 + \lambda x^2)^{\frac{1}{2}}}.$$

So, equation (3.35) becomes

$$K_t = 0$$

$$K = C_3(x).$$

Choosing $C_2(t) = 1$ and $C_3(x) = \frac{1}{(1 + \lambda x^2)^{\frac{1}{2}}}$, then

$$K = \frac{1}{(1 + \lambda x^2)^{\frac{1}{2}}}.$$

So, equation (3.33) becomes

$$F_t = -1$$

$$F = -t + C_4(x).$$

Choosing $C_4(x) = 0$, we have

$$F = -t.$$

So that, one obtains the linearizing transformation

$$X = -t, \quad dT = \frac{1}{(1 + \lambda x^2)^{\frac{1}{2}}} x' dt. \quad (3.39)$$

Hence, equation (3.32) is mapped by the transformation (3.39) into the linear equation

$$X'' = 0. \quad (3.40)$$

The general solution of equation (3.40) is

$$X = c_1 T + c_2, \quad (3.41)$$

where c_1 and c_2 , are arbitrary constants. Applying the generalized linearizing transformation (3.39) to equation (3.41), we obtain that the general solution of equation (3.32) is

$$-t = c_1 \phi(t) + c_2,$$

where the function $T = \phi(t)$ is a solution of the equation

$$\frac{dT}{dt} = \frac{1}{(1 + \lambda x^2)^{\frac{1}{2}}} x'.$$

3.5. Equation that can be linearizable by point and Sundman transformations

Consider the nonlinear second-order ordinary differential equation

$$x'' + \mu_3 x^{k_3} x'^2 + \mu_2 x^{k_2} x' + \mu_1 x^{k_1} = 0, \quad (3.42)$$

where $k_3, k_2, k_1, \mu_1, \mu_2$ and $\mu_3 \neq 0$ are arbitrary constants. The Lie criteria [4], showed that the nonlinear equation (3.42) is linearizable by a point transformation if and only if $\mu_1 = 0$ and $\mu_2 = 0$. In [4], Nakpim and Meleshko showed that the nonlinear equation (3.42) is linearizable by a generalized Sundman transformation if and only if $\mu_2 \neq 0$ and $\mu_1 = 0$.

By using our obtained theorems, we get the results as follow. Equation (3.42) is an equation of the form (2.4) in Theorem 2.1 with the coefficients

$$A_3 = 0, A_2 = \mu_3 x^{k_3}, A_1 = \mu_2 x^{k_2}, A_0 = \mu_1 x^{k_1}, \lambda_1 = k_2 \mu_2 x^{k_2},$$

$$\lambda_2 = x^{(k_1+k_3)} \mu_1 \mu_3 k_1 x + x^{(k_1+k_3)} \mu_1 \mu_3 k_3 x + x^{k_1} \mu_1 k_1^2 + x^{k_1} \mu_1 k_1 - x^{2k_2} \mu_2^2 k_2 x.$$

Now, the conditions in Theorem 2.2. case (a) is satisfied when the following conditions holds, that are,

$$k_2 \mu_2 x^{k_2} = 0,$$

$$x^{(k_1+k_3)} \mu_1 \mu_3 k_1 x + x^{(k_1+k_3)} \mu_1 \mu_3 k_3 x + x^{k_1} \mu_1 k_1^2 + x^{k_1} \mu_1 k_1 - x^{2k_2} \mu_2^2 k_2 x = 0.$$

Two cases arise.

Case 1: $\mu_2 = 0$ and $\mu_1 = 0$

In this case, the equation (3.42) takes the form

$$x'' + \mu_3 x^{k_3} x'^2 = 0. \quad (3.43)$$

The linearizing transformation is found by solving equations in Corollary 2.3 case (a), which become

$$F_t = -\frac{K}{G_1}, \quad (3.44)$$

$$K_x = \mu_3 x^{k_3} K, \quad (3.45)$$

$$K_t = \frac{K(2G_{1t} + \mu_3 x^{k_3} G_1 G_3)}{G_1}, \quad (3.46)$$

$$G_{3t} = \mu_3 x^{k_3} G_3^2, \quad (3.47)$$

$$G_{3xx} = -\frac{\mu_3 k_3 x^{k_3} G_3}{x}, \quad (3.48)$$

$$G_{1x} = \mu_3 x^{k_3} G_1. \quad (3.49)$$

Consider equation (3.49), we have

$$\frac{G_{1x}}{G_1} = \mu_3 x^{k_3}$$

$$\begin{aligned}\int \frac{1}{G_1} dG_1 &= \int \mu_3 x^{k_3} dx \\ \ln G_1 &= \frac{\mu_3 x^{k_3+1}}{k_3+1} + C(t) \\ G_1 &= C_1(t) e^{\frac{\mu_3 x^{k_3+1}}{k_3+1}}.\end{aligned}$$

Choosing $C_1(t) = 1$, one obtains

$$G_1 = e^{\frac{\mu_3 x^{k_3+1}}{k_3+1}}.$$

From equations (3.47) and (3.48), there exists $G_3 = 0$ such that these equations are hold. Then, let

$$G_3 = 0$$

so,

$$G_2 = 0.$$

Consider equation (3.45)

$$\begin{aligned}\frac{K_x}{K} &= \mu_3 x^{k_3} \\ \int \frac{1}{K} dK &= \int \mu_3 x^{k_3} dx \\ \ln K &= \frac{\mu_3 x^{k_3+1}}{k_3+1} + C(t) \\ K &= C_2(t) e^{\frac{\mu_3 x^{k_3+1}}{k_3+1}}.\end{aligned}$$

So, equation (3.46) becomes

$$\begin{aligned}K_t &= 0 \\ K &= C_3(x).\end{aligned}$$

Choosing $C_2(t) = 1$ and $C_3(x) = e^{\frac{\mu_3 x^{k_3+1}}{k_3+1}}$ then

$$K = e^{\frac{\mu_3 x^{k_3+1}}{k_3+1}}.$$

Thus, equation (3.44) becomes

$$\begin{aligned}F_t &= -1 \\ F &= -t + C_4(x).\end{aligned}$$

Choosing $C_4(x) = 0$, so

$$F = -t.$$

So that, one obtains the linearizing transformation

$$X = -t, \quad dT = e^{\frac{\mu_3 x^{k_3+1}}{k_3+1}} x' dt. \quad (3.50)$$

Hence, equation (3.43) is mapped by the transformation (3.50) into the linear equation

$$X'' = 0. \quad (3.51)$$

The general solution of equation (3.51) is

$$X = c_1 T + c_2, \quad (3.52)$$

where c_1 and c_2 , are arbitrary constants. Applying the generalized linearizing transformation (3.50) to equation (3.52), we obtain that the general solution of equation (3.43) is

$$-t = c_1 \phi(t) + c_2,$$

where the function $T = \phi(t)$ is a solution of the equation

$$\frac{dT}{dt} = e^{\frac{\mu_3 x^{k_3+1}}{k_3+1}} x',$$

where $k_3 \neq -1$.

For $k_3 = -1$.

Consider equation (3.49), we have

$$\begin{aligned} \frac{G_{1x}}{G_1} &= \frac{\mu_3}{x} \\ \int \frac{1}{G_1} dG_1 &= \int \frac{\mu_3}{x} dx \\ \ln G_1 &= \mu_3 \ln x + C(t) \\ G_1 &= C_5(t) x^{\mu_3}. \end{aligned}$$

Choosing $C_5(t) = 1$, one obtains

$$G_1 = x^{\mu_3}.$$

From equations (3.47) and (3.48), there exists $G_3 = 0$ such that these equations are hold. Then, let

$$G_3 = 0$$

so,

$$G_2 = 0.$$

Consider equation (3.45)

$$\begin{aligned} \frac{K_x}{K} &= \frac{\mu_3}{x} \\ \int \frac{1}{K} dK &= \int \frac{\mu_3}{x} dx \\ \ln K &= \mu_3 \ln x + C(t) \\ K &= C_6(t) x^{\mu_3}. \end{aligned}$$

อ. อ. 427
รศ. รศ. 5
รศ. 63

1049049



25

สำนักหอสมุด

28 ก.พ. 2565

So, equation (3.46) becomes

$$\begin{aligned} K_t &= 0 \\ K &= C_7(x). \end{aligned}$$

Choosing $C_6(t) = 1$ and $C_7(x) = x^{\mu_3}$ then

$$K = x^{\mu_3}.$$

Thus, equation (3.44) becomes

$$\begin{aligned} F_t &= -1 \\ F &= -t + C_8(x). \end{aligned}$$

Choosing $C_8(x) = 0$, so

$$F = -t.$$

So that, one obtains the linearizing transformation

$$X = -t, \quad dT = x^{\mu_3} x' dt. \quad (3.53)$$

Hence, equation (3.42) is mapped by the transformation (3.53) into the linear equation

$$X'' = 0. \quad (3.54)$$

The general solution of equation (3.54) is

$$X = c_1 T + c_2, \quad (3.55)$$

where c_1 and c_2 , are arbitrary constants. Applying the generalized linearizing transformation (3.53) to equation (3.55), we obtain that the general solution of equation (3.42) is

$$-t = c_1 \phi(t) + c_2,$$

where the function $T = \phi(t)$ is a solution of the equation

$$\frac{dT}{dt} = x^{\mu_3} x'.$$

Case 2: $k_2 = 0$ and $\mu_1 = 0$

In this case, the equation (3.12) takes the form

$$x'' + \mu_3 x^{k_3} x'^2 + \mu_2 x' = 0. \quad (3.56)$$

The linearizing transformation is found by solving equations in Corollary 2.3 case (a), which become

$$F_t = -\frac{K}{G_1}, \quad (3.57)$$

$$K_x = K \mu_3 x^{k_3}, \quad (3.58)$$

$$K_t = \frac{K(2G_{1t} + \mu_3 x^{k_3} G_1 G_3 - \mu_2 G_1)}{G_1}, \quad (3.59)$$

$$G_{3t} = G_3(x^{k_3} G_3 \mu_3 - \mu_2), \quad (3.60)$$

$$G_{3xx} = -\frac{G_3 \mu_3 k_3 x^{k_3}}{x}, \quad (3.61)$$

$$G_{1x} = G_1 \mu_3 x^{k_3}. \quad (3.62)$$

Consider equation (3.62), we have

$$\begin{aligned} \frac{G_{1x}}{G_1} &= \mu_3 x^{k_3} \\ \int \frac{1}{G_1} dG_1 &= \int \mu_3 x^{k_3} dx \\ \ln G_1 &= \frac{\mu_3 x^{k_3+1}}{k_3+1} + C(t) \\ G_1 &= C_1(t) e^{\frac{\mu_3 x^{k_3+1}}{k_3+1}}. \end{aligned}$$

Choosing $C_1(t) = 1$, one obtains

$$G_1 = e^{\frac{\mu_3 x^{k_3+1}}{k_3+1}}.$$

From equations (3.60) and (3.61), there exists $G_3 = 0$ such that these equations are hold. Then, let

$$G_3 = 0$$

so,

$$G_2 = 0.$$

Consider equation (3.58)

$$\begin{aligned} \frac{K_x}{K} &= \mu_3 x^{k_3} \\ \int \frac{1}{K} dK &= \int \mu_3 x^{k_3} dx \\ \ln K &= \frac{\mu_3 x^{k_3+1}}{k_3+1} + C(t) \\ K &= C_2(t) e^{\frac{\mu_3 x^{k_3+1}}{k_3+1}}. \end{aligned}$$

So, equation (3.59) becomes

$$\begin{aligned} \frac{K_t}{K} &= -\mu_2 \\ \int \frac{1}{K} dK &= -\mu_2 t + C(x) \\ K &= C_3(x) e^{-\mu_2 t}. \end{aligned}$$

Choosing $C_2(t) = e^{-\mu_2 t}$ and $C_3(x) = e^{\frac{\mu_3 x^{k_3+1}}{k_3+1}}$ then

$$K = e^{\frac{\mu_3 x^{k_3}}{k_3+1} - \mu_2 t}.$$

Thus, equation (3.57) becomes

$$\begin{aligned} F_t &= -e^{-\mu_2 t} \\ F &= \frac{e^{\mu_2 t}}{\mu_2} + C_4(x). \end{aligned}$$

Choosing $C_4(x) = 0$, so

$$F = \frac{e^{\mu_2 t}}{\mu_2}.$$

So that, one obtains the linearizing transformation

$$X = \frac{e^{\mu_2 t}}{\mu_2}, \quad dT = e^{\frac{\mu_3 x^{k_3+1}}{k_3+1}} x' dt. \quad (3.63)$$

Hence, equation (3.56) is mapped by the transformation (3.63) into the linear equation

$$X'' = 0. \quad (3.64)$$

The general solution of equation (3.64) is

$$X = c_1 T + c_2, \quad (3.65)$$

where c_1 and c_2 , are arbitrary constants. Applying the generalized linearizing transformation (3.63) to equation (3.65), we obtain that the general solution of equation (3.56) is

$$\frac{e^{\mu_2 t}}{\mu_2} = c_1 \phi(t) + c_2,$$

where the function $T = \phi(t)$ is a solution of the equation

$$\frac{dT}{dt} = e^{\frac{\mu_3 x^{k_3+1}}{k_3+1}} x',$$

where $k_3 \neq -1$.

For $k_3 = -1$.

Consider equation (3.62), we have

$$\begin{aligned} \frac{G_{1x}}{G_1} &= \frac{\mu_3}{x} \\ \int \frac{1}{G_1} dG_1 &= \int \frac{\mu_3}{x} dx \\ \ln G_1 &= \mu_3 \ln x + C(t) \end{aligned}$$

$$G_1 = C_5(t)x^{\mu_3}.$$

Choosing $C_5(t) = 1$, one obtains

$$G_1 = x^{\mu_3}.$$

From equations (3.60) and (3.61), there exists $G_3 = 0$ such that these equations are hold. Then, let

$$G_3 = 0$$

so,

$$G_2 = 0.$$

Consider equation (3.58)

$$\begin{aligned} \frac{K_x}{K} &= \frac{\mu_3}{x} \\ \int \frac{1}{K} dK &= \int \frac{\mu_3}{x} dx \\ \ln K &= \mu_3 \ln x + C(t) \\ K &= C_6(t)x^{\mu_3}. \end{aligned}$$

So, equation (3.59) becomes

$$\begin{aligned} \frac{K_t}{K} &= -\mu_2 \\ \int \frac{1}{K} dK &= -\mu_2 t + C(x) \\ K &= C_7(x)e^{-\mu_2 t}. \end{aligned}$$

Choosing $C_6(t) = e^{-\mu_2 t}$ and $C_7(x) = x^{\mu_3}$ then

$$K = x^{\mu_3} e^{-\mu_2 t}.$$

Thus, equation (3.57) becomes

$$\begin{aligned} F_t &= -e^{\mu_2 t} \\ F &= \frac{e^{\mu_2 t}}{\mu_2} + C_8(x). \end{aligned}$$

Choosing $C_8(x) = 0$, so

$$F = \frac{e^{\mu_2 t}}{\mu_2}.$$

So that, one obtains the linearizing transformation

$$X = \frac{e^{\mu_2 t}}{\mu_2}, \quad dT = x^{\mu_3} x' dt. \quad (3.66)$$

Hence, equation (3.56) is mapped by the transformation (3.66) into the linear equation

$$X'' = 0. \quad (3.67)$$

The general solution of equation (3.67) is

$$X = c_1 T + c_2, \quad (3.68)$$

where c_1 and c_2 , are arbitrary constants. Applying the generalized linearizing transformation (3.66) to equation (3.68), we obtain that the general solution of equation (3.56) is

$$\frac{e^{\mu_2 t}}{\mu_2} = c_1 \phi(t) + c_2,$$

where the function $T = \phi(t)$ is a solution of the equation

$$\frac{dT}{dt} = x^{\mu_3} x',$$

Remark:

From theorem 2.2. case (b) equation (3.42) is linearizable if only if $\mu_1 = 0$.

3.6. Modified generalized Vakhnenko equation

In 2009, Ma, Li and Wang [4] considered a modified generalized Vakhnenko equation (mGVE),

$$\frac{\partial}{\partial x} (\mathfrak{D}^2 u + \frac{1}{2} \rho u^2 + \beta u) = 0, \quad \mathfrak{D} = \frac{\partial}{\partial t} + u \frac{\partial}{\partial x}, \quad (3.69)$$

where ρ , q , β are arbitrary non-zero constants.

To construct the exact solutions for mGVE is all important. Many studies have been conducted. For examples, when $\rho = \beta = 0$ and $q = 1$, equation (3.69) is reduced to well-known Vakhnenko equation (VE), which governs the nonlinear propagation of high-frequency wave in a relaxing medium [4]-[4]. The VE has soliton solutions [4]. When $\rho = q = 1$ and β an arbitrary non-zero constant, equation (3.69) is reduced as the generalized VE (GVE), in [4] it was shown that GVE has N-soliton solution. When $\rho = 2q$ and β an arbitrary non-zero constant, equation (3.69) has a loop-like, hump-like and cusp-like soliton solutions [4]. In [4], it was shown that equation (3.69) has travelling wave solution and single-soliton solution.

Consider a modified generalized Vakhnenko equation (3.69), we can rewrite it in the form

$$2u_t u_{tx} + 2[uu_x u_{tx} + u_t (uu_{xx} + u_x^2)] + 2u^2 u_{xx} + 2uu_x^3 + \rho uu_x + \beta u_x + q(u_t + uu_x) = 0. \quad (3.70)$$

Of particular interest among solutions of equation (3.70) are travelling wave solutions:

$$u(t, x) = H(x - Dt),$$

where D is a constant phase velocity and the argument $x-Dt$ is a phase of the wave. Substituting the representation of a solution into equation (3.70), one finds

$$2D^2H'H'' - 2DH'(2HH'' + H'^2) + 2H^2H'H'' + 2HH'^3 + \rho HH' + \beta H' + q(-DH' + HH') = 0. \quad (3.71)$$

By using the obtained theorems, we get the results as follow. Equation (3.71) is an equation of the form in Theorem 2.1 with the coefficients

$$A_3 = 0, A_2 = -\frac{1}{(D-H)}, A_1 = 0, A_0 = \frac{\rho H + \beta - qD + qH}{2(D^2 - 2DH + H^2)},$$

$$\lambda_1 = 0, \lambda_2 = \rho D + \beta.$$

From Theorem 2.2. case (a), equation (3.71) is linearizable if only if $\rho D + \beta = 0$.

3.7. Burgers' equation

Burgers' equation is obtained as a result of combining nonlinear wave motion with linear diffusion and is the simplest model for analyzing combined effect of nonlinear advection and diffusion. The presence of viscous term helps suppress the wave-breaking, smooth out shock discontinuities, and hence we expect to obtain a well-behaved and smooth solution. Moreover, in the inviscid limit, as the diffusion term becomes vanishingly small, the smooth viscous solutions converge non-uniformly to the appropriate discontinuous shock wave, leading to an alternative mechanism for analyzing conservative nonlinear dynamical processes. In 2016, A. Salih [4] considered the nonlinear convection-diffusion equation

$$\frac{\partial u}{\partial t} + u \frac{\partial u}{\partial x} - v \frac{\partial^2 u}{\partial x^2} = 0, v > 0 \quad (3.72)$$

which is known as Burgers' equation. This equation is balance between time evolution, nonlinearity, and diffusion. This is the simplest nonlinear model equation for diffusive waves in fluid dynamics. Burgers (1948) first developed this equation primarily to throw light on turbulence described by the interaction of two opposite effects of convection and diffusion.

The term uu_x will have a shocking up effect that will cause waves to break and the term vu_{xx} is a diffusion term like the one occurring in the heat equation.

Of particular interest among solutions of equation (3.72) are travelling wave solutions:

$$u(t, x) = H(x - Dt),$$

where D is a constant phase velocity and the argument $x-Dt$ is a phase of the wave. Substituting the representation of a solution into equation (3.72), one finds

$$-DH' + HH' - vH'' = 0. \quad (3.73)$$

By using the obtained theorems, we get the results as follow. Equation (3.73) is an equation of the form in Theorem 2.1 with the coefficients

$$A_3 = 0, A_2 = 0, A_1 = \frac{D-H}{v}, A_0 = 0, \lambda_1 = \frac{1}{v}, \lambda_2 = \frac{-D+H}{v^2}.$$

One can check that these coefficients obey the condition in Theorem 2.2. case (b). Thus, equation (3.73) is linearizable via a generalized linearizing transformation.

4. Conclusion

In this research, the necessary condition which guarantee that the second-order ordinary differential equation can be linearized by generalized linearizing transformation is found in Theorem 2.1. Theorem 2.2 case (a) and case (b) are sufficient conditions for the linearization problem, they are selected by the value of λ_1 . A new algorithm for finding linearizing transformation is summarized in Corollary 2.3. Finally, some applications are provided to demonstrate our procedure.

References

- [1] S. V. Meleshko, *Methods for Constructing Exact Solutions of Partial Differential Equations*, Springer Science+Business Media, Inc., New York, 2005, mathematical and Analytical Techniques with Applications to Engineering.
- [2] N. H. Ibragimov, *Elementary Lie Group Analysis and Ordinary Differential Equations*, Wiley & Sons, Chichester, 1999.
- [3] L. G. S. Durate, I. C. Moreira, F. C. Santos, Linearisation under nonpoint transformations, *J Phys A : Math Gen* 27 (1994) L739-L743.
- [4] J. R. C. Muriel, Nonlocal transformations and linearization of second-order ordinary differential equations, *J. Phys. A: Math. Theor.* 43 (434025) (2010) 13 pages.
- [5] A. A.-D. M.T. Mustafa, R. Mara'beh, On the linearization of second-order ordinary differential equations to the laguerre form via generalized sundman transformations, *SIGMA* 9 (041) (2013) 10 pages.
- [6] W. Nakpim, S. Meleshko, Linearization of third-order ordinary differential equations by generalized sundman transformations: the case $X''' + \alpha X = 0$, *Communications in Nonlinear Science and Numerical Simulation* 15 (2010) 1717-1723.
- [7] V. K. Chandrasekar, Senthilvelan, M. Lakshmanan, A unification in the theory of linearization of second-order nonlinear ordinary differential equations, *Journal of Physics A: Mathematical and General* 39 (2006) L69-L76.
- [8] V. K. Chandrasekar, M. Lakshmanan, On the complete integrability and linearization of nonlinear ordinary differential equations, *Proc. R. Soc. A* 461 (2060) (2005) 2451-2477.
- [9] V. K. Chandrasekar, M. Lakshmanan, On the complete integrability and linearization of nonlinear ordinary differential equations. II. Third-order equations, *Proc. R. Soc. A* 462 (2006) 1831-1852.
- [10] S. Lie, (1883). Klassifikation und Integration von gewöhnlichen Differentialgleichungen zwischen x, y , die eine Gruppe von Transformationen gestatten. III Archiv for Matematik og Naturvidenskab, 8(4), 371-427. Reprinted in Lie's Gessammelte Abhandlungden, 1924, 5, paper XIY, pp. 362-427.
- [11] <https://math.stackexchange.com/questions/397461/examples-of-nonlinear-ordinary-differential-equations-with-elementary-solutions>.
- [12] L. H. Koudahoun, J. Akande, D. K. K. Adjai, Y. J. F. Kpomahou, M. D. Monsia, Exact Classical and Quantum Mechanics of a Generalized Singular Equation of Quadratic Liénard Type, *Journal of Mathematics and Statistics*. (2018), Volume 14: 183.192.
- [13] M. T. Mustafa, A. Y. Al-Dweik, R. A. Mara'beh, On the linearization of second-order ordinary differential equations to the laguerre form via generalized Sundman transformations, *Symmetry, Integrability and Geometry: Methods and Applications (SIGMA)*; vol. 9, article no. 041 (2013).
- [14] P. M. Mathews, M. Lakshmanan, On a unique nonlinear oscillator, *Q. Appl. Math.*, 32, 215-218 (1974).
- [15] Nakpim W., Meleshko S.V., Linearization of second-order ordinary differential equations by generalized Sundman transformations, *SIGMA* 6 (2010), 051, 11 pages.
- [16] Y. Ma, B. Li, C. Wang, A series of abundant exact travelling wave solutions for a modified generalized Vakhnenko equation using auxiliary equation method, *Applied Mathematics and Computation*, 211, 102-107 (2009).
- [17] V.O. Vakhnenko, E.J. Parkes, The two loop soliton solution of the Vakhnenko equation, *Nonlinearity*, 11, 1457-1464 (1998).
- [18] V.O. Vakhnenko, High-frequency soliton-like waves in a relaxing medium, *J. Math. Phys.*, 40, 2011-2020 (1999).
- [19] V.O. Vakhnenko, E.J. Parkes, The calculation of multi-soliton solutions of the Vakhnenko equation by the inverse scattering method, *Chaos Solitons Fract*, 13, 1819-1826 (2002).

- [20] A.J. Morrison, E.J. Parkes, The N-soliton solution of the modified generalized Vakhnenko equation (a new nonlinear evolution equation), *Chaos Solitons Fract*, 16, 13–26 (2003).
- [21] V.O. Vakhnenko, E.J. Parkes, A Bäcklund transformation and the inverse scattering transform method for the generalised Vakhnenko equation, *Chaos Solitons Fract*, 17, 683–692 (2003).
- [22] Y.P. Liu, Z.B. Li, K.C. Wang, Symbolic computation of exact solutions for a nonlinear evolution equation, *Chaos Solitons Fract*, 31, 1173–1180 (2007).
- [23] A. Salih, *Burgers' Equation*, (2016).



ภาคผนวก

ประกอบด้วย

- บทความสำหรับการเผยแพร่
- ตารางเปรียบเทียบกิจกรรมที่วางแผนไว้ และกิจกรรมที่ดำเนินการมา
- ผลที่ได้รับตลอดโครงการ



บทความสำหรับการเผยแพร่

ได้ตีพิมพ์ผลงานในวารสารวิชาการระดับนานาชาติในฐานข้อมูลของ SJR อยู่ใน Quartile 3 จำนวน 1 เรื่อง ได้แก่

Prakrong Voraka, Supaporn Suksern, and Nontakan Donjiwprai. Linearizability of Nonlinear Second-Order Ordinary Differential Equations by Using a Generalized Linearizing Transformation. IAENG International Journal of Applied Mathematics., 50:4, IJAM_50_4_19



Linearizability of Nonlinear Second-Order Ordinary Differential Equations by Using a Generalized Linearizing Transformation

Prakrong Voraka, Supaporn Suksern, and Nontakan Donjiwprai

Abstract—In this paper, we have proposed the linearization problem of second-order ordinary differential equation under the generalized linearizing transformation. We found the necessary form for reducing the second-order ordinary differential equation to simple linear equation. We also obtained sufficient condition for making the above form to be linear. Further, the procedure of linear transformation within the study is demonstrated in the explicit form. Moreover, we apply the obtained linearization criteria to the interesting problems of nonlinear ordinary differential equations and nonlinear partial differential equations, for examples the parachute equation, the Painlevé - Gambier XI equation, the equation for the variable frequency oscillator, the one-dimensional non-polynomial oscillator, the equation that can be linearizable by point and Sundman transformations, the modified generalized Vakhnenko equation.

Index Terms—linearization problem, generalized linearizing transformation, nonlinear second-order ordinary differential equation.

I. Introduction

THE linearization problem is one of the important branches in differential equation field. A number of mathematicians has been studying this branch continuously until the present time. To discover theory for finding new knowledge has shown to be a great benefit for academic world and country development. It is known that theories and new knowledge obtained from research not only offer benefits to improve existing knowledge within the branch itself, but also they can be applied to other branches or fields and can be key fundamental to develop basic science which is basic research to build many other new knowledge. This would be a fundamental step to develop the country.

The linearization problem is a branch of study that can be applied widely in particular to the study involving solving the equations. Most important physical problems are in the form of nonlinear differential equations which are normally difficult to solve and there are relatively few method to find their exact solutions. Numerical method therefore is often used to solve these nonlinear

differential equations but the obtained solutions are just the approximate solutions. However, the exact solution is claimed to be more interesting because it can be used to analyze the properties of the studied equations. One of the methods used to determine the exact solutions is to linearize the interested equation and find solutions directly by fundamental method. The solutions obtained from such linear equation are yet still solutions of initial equation. By mentioned above, we are required to seek for transformation in order to transform initial equation to be linear equation.

There are a number of interesting transformations. For example, in the case that the transformation consists of derivative, we call it as tangent transformation, in the case that the transformation depends only on independent and dependent variables, we call it as point transformation and we will call the tangent transformation which the independent and dependent variables can be changed and involves the first derivative as contact transformation. In addition, another type of transformation which its transformation set is different from any mentioned above since there is a nonlocal term $T = \int G(t, x) dt$, such transformation is called generalized Sundman transformation. In this paper, we use the generalized linearizing transformation which is an extended transformation from generalized Sundman transformation where the selected G function is $G(t, x, x')$.

Up to the present time, all researchers who study the linearization of second-order ordinary differential equations via generalized linearizing transformation have not covered all cases yet. Therefore, in this paper we focus on the remaining cases that have not yet been studied, which we also find that those cases can be applied to solve several nonlinear equations in real-world phenomenon.

A. Historical Review

From above facts as mentioned, the researcher would like to give a brief background of this study. Since 19th century the linearization problem of ordinary differential equation has attracted some interests from various well-known mathematicians e.g. S. Lie and E. Cartan etc. The first person who could solve the linearization problem of ordinary differential equation is Lie [1]. Lie could discover the standard form of every second-order ordinary differential equation which could be reduced the form to become linear equation via changing the

Manuscript received June 11, 2020; revised August 17, 2020. This work was financially supported by Naresuan University, Thailand under Grant no. R2563C017.

S. Suksern and N. Donjiwprai (corresponding author e-mail: supapornsu@nu.ac.th) are with the Department of Mathematics, Faculty of Science, Naresuan University, Phitsanulok 65000, Thailand.

S. Suksern is with the Research center for Academic Excellence in Mathematics, Naresuan University, Phitsanulok 65000, Thailand.

P. Voraka is with the Faculty of Science and Engineering, Kasetsart University, Chalermphrakiat Sakon Nakhon Province Campus, Sakon Nakhon 47000, Thailand.

independent and dependent variables (or can be called point transformation). Later, Liouville [2] and Tresse [3] used the relative invariants of equivalence group under point transformation to solve the equivalence of second-order ordinary differential equations which can be reduced from second-order nonlinear ordinary differential equations to second-order linear ordinary differential equations. Moreover, Lie discovered that every second-order ordinary differential equation can be reduced to second-order linear ordinary differential equation without any conditions via contact transformation.

Having mentioned some methods above, there are yet still other methods to solve linearization problem of second-order ordinary differential equation. For example, the method of Cartan [4], the reducing order method, the differential substitution method etc.

Another transformation that is very interesting and has not been mentioned yet is the generalized Sundman transformation

$$X = F(t, x), \quad dT = G(t, x)dt. \quad (1)$$

Duarte, Moreira and Santos [5] used generalized Sundman transformation to determine the conditions for linearizing the second-order ordinary differential equation to be simple linear equation. In [6] Nakpim and Meleshko demonstrated that the general linear equation in the canonical form of Laguerre was not sufficient for solving linearization problem via generalized Sundman transformation. The canonical form of Laguerre could only particularly be applied with point and contact transformations. Therefore, in [6] they found the conditions for linearizing the second-order ordinary differential equation to be general linear equation.

In this paper, we extend the generalized Sundman transformation which was studied before as shown in [7]-[9], where they called such a transformation in this form as generalized linearizing transformation

$$X = F(t, x), \quad dT = G(t, x, x')dt. \quad (2)$$

They demonstrated that this transformation can be used to linearize a more extensive class of nonlinear standard differential equations including some equations that can't be linearized by the non-point and invertible point transformations. In the case that the function G in (2) does not depend on the variable x' , then it can be turned into a non-point transformation. If G is a differentiable function, then it turns into an invertible point transformation. In this way, (2) is a unified transformation as it incorporates non-point and invertible point transformations as extraordinary cases. A case of an equation that can be linearized by a change of the structure (2) is given in [8].

In [7], Chandrasekar, Senthilvelan and Lakshmanan applied a particular class of transformations (2), where the function $G(t, x, x')$ is linear with respect to x' .

They payed attention to the case where G is a polynomial function in x' and in particular where it is linear in x' with coefficients which are arbitrary functions of t and x . To be specific, they focused here on the case

$$X = F(t, x), \quad dT = (G_1(t, x) x' + G_2(t, x)) dt.$$

Notice that for the case $G_1 = 0$, the generalized linearizing transformation becomes a generalized Sundman transformation, so that they assumed $G_1 \neq 0$.

The authors of [7] obtained that any second-order linearizable ordinary differential equation which can be mapped into the equation $X'' = 0$ via a generalized linearizing transformation has to be of the form

$$x'' + A_3(t, x)x'^3 + A_2(t, x)x'^2 + A_1(t, x)x' + A_0(t, x) = 0, \quad (3)$$

and the functions A_i 's ($i = 0, 1, 2, 3$) are connected to the transform functions F and G through the relations

$$\begin{aligned} A_3 &= (G_1 F_{xx} - F_x G_{1x})/M, \\ A_2 &= (G_2 F_{xx} + 2G_1 F_{xt} - F_x G_{2x} - F_t G_{1x} - F_x G_{1t})/M, \\ A_1 &= (2G_2 F_{xt} + G_1 F_{tt} - F_x G_{2t} - F_t G_{2x} - F_t G_{1t})/M, \\ A_0 &= (G_2 F_{tt} - F_t G_{2t})/M \end{aligned} \quad (4)$$

with $M = F_x G_2 - F_t G_1 \neq 0$.

They have analyzed a particular case of equation (3), namely, $A_3 = 0$ and $A_2 = 0$ in equation (4). Complete analysis of the compatibility of arising equations is given for the case $F_x \neq 0$.

Therefore, in this paper we will apply the generalized linearizing transformation with second-order ordinary differential equation to complete the remaining cases ($F_x = 0$) which are different from the work by Chandrasekar and Lakshmanan [7].

II. Formulation of the Linearization Theorems

A. Obtaining Necessary Condition of Linearization

We begin with investigating the necessary conditions for linearization. We consider the second-order ordinary differential equation

$$x'' = F(t, x, x') \quad (5)$$

which can be transformed to a simplest linear equation

$$X'' = 0 \quad (6)$$

under the generalized linearizing transformation

$$\begin{aligned} X &= F(t, x), \\ dT &= [G_1(t, x) x' + G_2(t, x)] dt, \end{aligned} \quad (7)$$

where $G_1 \neq 0$. So, we arrive at the following theorem.

Theorem 2.1: Any second-order ordinary differential equations (5) obtained from a linear equation (6) by a generalized linearizing transformation (7) has to be the form

$$x'' + A_3(t, x)x'^3 + A_2(t, x)x'^2 + A_1(t, x)x' + A_0(t, x) = 0, \quad (8)$$

where

$$A_3 = (-F_{xx}G_1 + F_x G_{1x})/(F_t G_1 - F_x G_2), \quad (9)$$

$$A_2 = (-2F_{tx}G_1 + F_t G_{1x} - F_{xx}G_2 + F_x G_{1t} + F_x G_{2x})/(F_t G_1 - F_x G_2), \quad (10)$$

$$A_1 = (-2F_{tx}G_2 - F_{tt}G_1 + F_t G_{1t} + F_t G_{2x} + F_x G_{2t})/(F_t G_1 - F_x G_2), \quad (11)$$

$$A_0 = (-F_{tt}G_2 + F_t G_{2t})/(F_t G_1 - F_x G_2). \quad (12)$$

Proof. Applying a generalized linearizing transformation (7), one obtains the following transformations

$$X'(T) = \frac{D_t F}{D_t \int [G_1 x' + G_2] dt} = \frac{F_t + x' F_x}{G_1 x' + G_2} = P(t, x, x'),$$

$$X''(T) = \frac{D_t P}{D_t \int [G_1 x' + G_2] dt} = \frac{P_t + P_x x' + P_{x'} x''}{G_1 x' + G_2},$$

where

$$P_t = \frac{F_{tt}(G_1 x' + G_2) - F_t(G_1 x'' + G_{1t} x' + G_{2t})}{(G_1 x' + G_2)^2},$$

$$P_x = \frac{F_{tx}(G_1 x' + G_2) - F_t(G_{1x} x' + G_{2x})}{(G_1 x' + G_2)^2},$$

$$P_{x'} = -\frac{F_t G_{1x'}}{(G_1 x' + G_2)^2},$$

and $D_t = \frac{\partial}{\partial t} + x' \frac{\partial}{\partial x} + x'' \frac{\partial}{\partial x'} + \dots$ is a total derivative. Substituting the resulting expression into the linear equation (6) we arrive at the necessary form (8), where A_0, A_1, A_2 and A_3 are some functions of t and x as defined in system of equations (9)-(12).

B. Obtaining Sufficient Conditions of Linearization and Linearizing Transformation

For obtaining sufficient conditions of linearizability of equation (8), one has to solve the compatibility problem of the system of equations (9)-(12), considering it as overdetermined system of partial differential equations for the functions F, G_1 and G_2 with given coefficients A_i of equation (8).

For convenience of calculations, we set

$$G_3 = \frac{G_2}{G_1}.$$

So that system of equations (9)-(12) become

$$A_3 = (-F_{xx} G_1 + F_x G_{1x}) / (G_1 (F_t - F_x G_3)), \quad (13)$$

$$A_2 = (-2F_{tx} G_1 + F_t G_{1x} - F_{xx} G_1 G_3 + F_x G_{1t} + F_x G_{1x} G_3 + F_x G_{3x} G_1) / (G_1 (F_t - F_x G_3)), \quad (14)$$

$$A_1 = (-2F_{tx} G_1 G_3 - F_{tt} G_1 + F_t G_{1t} + F_t G_{1x} G_3 + F_t G_{3x} G_1 + F_x G_{1t} G_3 + F_x G_{3t} G_1) / (G_1 (F_t - F_x G_3)), \quad (15)$$

$$A_0 = (-F_{tt} G_1 G_3 + F_t G_{1t} G_3 + F_t G_{3t} G_1) / (G_1 (F_t - F_x G_3)). \quad (16)$$

According to the notation $K = G_1 (F_x G_3 - F_t)$, we define the derivative F_t as

$$F_t = (F_x G_1 G_3 - K) / G_1. \quad (17)$$

Solving equations (13)-(16) with respect to F_{xx}, K_x, K_t

and G_{3t} , one finds

$$F_{xx} = (F_x G_{1x} + A_3 K) / G_1, \quad (18)$$

$$K_x = (-F_x G_{1t} G_1 + F_x G_{1x} G_1 G_3 + F_x G_{3x} G_1^2 + 3G_{1x} K - A_2 G_1 K + 3A_3 G_1 G_3 K) / (2G_1), \quad (19)$$

$$K_t = (-F_x G_{1t} G_1 G_3 + F_x G_{1x} G_1 G_3^2 + F_x G_{3x} G_1^2 G_3 + 4G_{1t} K - G_{1x} G_3 K + 2G_{3x} G_1 K - 2A_1 G_1 K + 3A_2 G_1 G_3 K - 3A_3 G_1 G_3^2 K) / (2G_1), \quad (20)$$

$$G_{3t} = G_{3x} G_3 + A_0 - A_1 G_3 + A_2 G_3^2 - A_3 G_3^3. \quad (21)$$

Comparing the mixed derivative $(K_x)_t = (K_t)_x$, one obtains

$$G_{3xx} = (2A_0 F_x G_1^3 - 2A_{1x} F_x G_1^3 G_3 + 4A_{1x} G_1^2 K - 2A_{2t} G_1^2 K + 2A_{2x} F_x G_1^3 G_3^2 - 6A_{2x} G_1^2 G_3 K + 6A_{3t} G_1^2 G_3 K - 2A_{3x} F_x G_1^3 G_3^2 + 6A_{3x} G_1^2 G_3^2 K + 4F_x G_{1tx} G_1^2 G_3 - 2F_x G_{1tt} G_1^2 + 3F_x G_{1t}^2 G_1 - 6F_x G_{1t} G_{1x} G_1 G_3 + 2F_x G_{1t} G_{3x} G_1^2 - 2F_x G_{1t} A_1 G_1^2 + 4F_x G_{1t} A_2 G_1^2 G_3 - 6F_x G_{1t} A_3 G_1^2 G_3^2 - 2F_x G_{1xx} G_1^2 G_3^2 + 3F_x G_{1x}^2 G_1 G_3^2 - 2F_x G_{1x} G_{3x} G_1^2 G_3 + 2F_x G_{1x} A_0 G_1^2 - 2F_x G_{1x} A_2 G_1^2 G_3^2 + 4F_x G_{1x} A_3 G_1^2 G_3^2 - F_x G_{3xx} G_1^3 - 2G_{1tx} G_1 K + 3G_{1t} G_{1x} K - G_{1t} A_2 G_1 K + 3G_{1t} A_3 G_1 G_3 K + 2G_{1xx} G_1 G_3 K - 3G_{1x}^2 G_3 K + G_{1x} G_{3x} G_1 K + G_{1x} A_2 G_1 G_3 K - 3G_{1x} A_3 G_1 G_3^2 K - 5G_{3xx} A_2 G_1^2 K + 15G_{3xx} A_3 G_1^2 G_3 K + 6A_0 A_3 G_1^2 K - 6A_1 A_3 G_1^2 G_3 K + 6A_2 A_3 G_1^2 G_3^2 K - 6A_3^2 G_1^2 G_3^2 K) / (4G_1^2 K). \quad (22)$$

The compatibility analysis depends on the value of F_x . A complete study of all cases is cumbersome. Here a complete solution is given for the case where $F_x = 0$.

Case $F_x = 0$

Since $F_x = 0$, then substituting it into F_{xx} in equation (18), one gets the condition

$$A_3 = 0. \quad (23)$$

Comparing the mixed derivative $(F_t)_x = (F_x)_t$, one obtains the derivative

$$G_{1x} = A_2 G_1 - 3A_3 G_1 G_3 \quad (24)$$

and this satisfies equation $(F_{xx})_t = (F_t)_{xx}$. Setting

$$\lambda_1 = -A_{1x} + 2A_{2t},$$

$$\lambda_2 = -A_{0xx} - A_{0x} A_2 + A_{2tt} + A_{2t} A_1 - A_{2x} A_0 - \lambda_{1t} - A_1 \lambda_1$$

then, equation $(G_{3xx})_t = (G_{3t})_{xx}$ becomes

$$G_{3xx} \lambda_1 + G_3 A_2 \lambda_1 + \lambda_2 = 0. \quad (25)$$

The compatibility analysis depends on the value of λ_1 . A complete study of all cases is given here.

3.3.1. Case $\lambda_1 = 0$

From equation (25), one finds the condition

$$\lambda_2 = 0. \tag{26}$$

3.3.2. Case $\lambda_1 \neq 0$

Equation (25) provides the derivative

$$G_{3x} = -(G_3 A_2 \lambda_1 + \lambda_2) / \lambda_1. \tag{27}$$

Substituting G_{3x} into G_{3xx} in equation (22), one arrives at the condition

$$\lambda_{2x} = (-A_{2t} \lambda_1^2 + \lambda_{1x} \lambda_2 + \lambda_1^3) / \lambda_1. \tag{28}$$

Comparing the mixed derivatives $(G_{3x})_t = (G_{3t})_x$, one gets the condition

$$\lambda_{2t} = -(A_{0x} \lambda_1^2 - \lambda_{1t} \lambda_2 + A_0 A_2 \lambda_1^2 + A_1 \lambda_1 \lambda_2 + \lambda_2^2) / \lambda_1. \tag{29}$$

Combining all derived results in the case $F_x = 0$ the following theorems are proven.

Theorem 2.2: Sufficient conditions for equation (8) to be equivalent to a linear equation (6) via generalized linearizing transformation (7) with the function $F = F(t)$ is the equation (23) and the additional conditions are as follows.

(a) If $\lambda_1 = 0$, then the condition is equation (26).

(b) If $\lambda_1 \neq 0$, then the conditions are equations (28) and (29).

Theorem 2.3: Provided that the sufficient conditions in Theorem 2.2 are satisfied, the transformation (7) with the function $F = F(t)$ mapping equation (8) to a linear equation (6) is obtained by solving the compatible system of equations :

(a) (17), (19), (20), (21), (22), and (24).

(b) (17), (19), (20), (21), (24), and (27).

III. Some Applications

In this section we focus on finding some applications which satisfy Theorem 2.1, Theorem 2.2 and Theorem 2.3. The obtained results are as follows.

A. Parachute Equation

An application to this equation can be applied to a model of motion for a parachutist by using Newton's law II which is $\sum F = ma$. The movement of skydiver when the coefficient of air opposition changes between free-fall and the last consistent state drop with the parachute is slowly conveyed.

Consider the parachute equation [10], in the form

$$x'' - kx'^2 + g = 0, \tag{30}$$

with initial conditions $x(0) = 0$ and $x'(0) = 0$.

Here $k = \frac{\pi \rho C_d D^2}{8m}$, where

- m is the mass of the body and parachute,
- ρ is the density of the fluid in which the body moves,
- C_d is the drag coefficient for the parachute (1.5 for parabolic profile and 0.75 for flat),
- D is the effective diameter of the parachute.

Equation (30) is an equation of the form (8) in Theorem 2.1 with the coefficients

$$A_3 = 0, A_2 = k, A_1 = 0, A_0 = g, \lambda_1 = 0, \lambda_2 = 0.$$

One can check that these coefficients obey the conditions in Theorem 2.2. case (a). Thus, equation (30) is linearizable via a generalized linearizing transformation. For finding the functions F , G_1 and G_2 we have to solve equations in Theorem 2.3 case (a), which become

$$F_t = -\frac{K}{G_1}, K_x = kK, K_t = \frac{K(2G_{1t} + G_1 G_3 k)}{G_1}, \tag{31}$$

$$G_{3t} = g + G_3^2 k, G_{3xx} = 0, G_{1x} = G_1 k.$$

One can find the particular solution for equations in (31) as

$$G_1 = e^{kx}, G_3 = \sqrt{\frac{g}{k}} i, G_2 = \sqrt{\frac{g}{k}} i e^{kx},$$

$$K = e^{kx + \sqrt{kg} it}, F = \frac{i}{\sqrt{kg}} e^{\sqrt{kg} it}.$$

So that, one obtains the linearizing transformation

$$X = \frac{i}{\sqrt{kg}} e^{\sqrt{kg} it}, dT = (e^{kx} x' + \sqrt{\frac{g}{k}} i e^{kx}) dt. \tag{32}$$

Hence, equation (30) is mapped by the transformation (32) into the linear equation

$$X'' = 0. \tag{33}$$

The general solution of equation (33) is

$$X = c_1 T + c_2, \tag{34}$$

where c_1 and c_2 are arbitrary constants. Applying the generalized linearizing transformation (32) to equation (34), we obtain that the general solution of equation (30) is

$$\frac{i}{\sqrt{kg}} e^{\sqrt{kg} it} = c_1 \phi(t) + c_2,$$

where the function $T = \phi(t)$ is a solution of the equation

$$\frac{dT}{dt} = (x' + \sqrt{\frac{g}{k}} i) e^{kx}.$$

B. Painlevé - Gambier XI Equation

In [11], Koudahoun, Akande, Adjai, Kpomahou and Monsia considered the Painlevé - Gambier XI equation

$$x'' + \frac{x'^2}{x} = 0. \tag{35}$$

To investigate the exact classical and quantum mechanical solutions, they offered a generalized singular differential equation of quadratic Lienard type.

By using our obtained theorems, we get the results as follow. Equation (35) is an equation of the form (8) in Theorem 2.1 with the coefficients

$$A_3 = 0, A_2 = \frac{1}{x}, A_1 = 0, A_0 = 0, \lambda_1 = 0, \lambda_2 = 0.$$

One can check that these coefficients obey the conditions in Theorem 2.2. case (a). Thus, equation (35) is linearizable via a generalized linearizing transformation. For finding the functions F , G_1 and G_2 we have to solve equations in Theorem 2.3 case (a), which become

$$F_t = -\frac{K}{G_1}, K_x = \frac{K}{x}, K_t = \frac{K(2G_{1t} x + G_1 G_3)}{G_1 x}, \tag{36}$$

$$G_{3t} = \frac{G_3^2}{x}, G_{3xx} = \frac{G_3}{x^2}, G_{1x} = \frac{G_1}{x}.$$

One can find the particular solution for equations in (36) as

$$G_1 = x, G_3 = 0, G_2 = 0, K = x, F = -t.$$

So that, one obtains the linearizing transformation

$$X = -t, \quad dT = xx' dt. \quad (37)$$

Hence, equation (35) is mapped by the transformation (37) into the linear equation

$$X'' = 0. \quad (38)$$

The general solution of equation (38) is

$$X = c_1 T + c_2, \quad (39)$$

where c_1 and c_2 are arbitrary constants. Applying the generalized linearizing transformation (37) to equation (39), we obtain that the general solution of equation (35) is

$$-t = c_1 \phi(t) + c_2,$$

where the function $T = \phi(t)$ is a solution of the equation

$$\frac{dT}{dt} = xx'.$$

C. Equation for the Variable Frequency Oscillator

In 2013, Mastafa, Al-Dueik and Mara'beh [12] considered the ordinary differential for the variable frequency oscillator

$$x'' + xx'^2 = 0. \quad (40)$$

They showed that this equation can be linearizable by generalized Sundman transformation.

By using our obtained theorems, we get the results as follow. Equation (40) is an equation of the form (8) in Theorem 2.1 with the coefficients

$$A_3 = 0, \quad A_2 = x, \quad A_1 = 0, \quad A_0 = 0, \quad \lambda_1 = 0, \quad \lambda_2 = 0.$$

One can check that these coefficients obey the conditions in Theorem 2.2. case (a). Thus, equation (40) is linearizable via a generalized linearizing transformation. For finding the functions F , G_1 and G_2 we have to solve equations in Theorem 2.3 case (a), which become

$$F_t = -\frac{K}{G_1}, \quad K_x = xK, \quad K_t = \frac{K(2G_{1t} + G_1 G_{3x})}{G_1}, \quad (41)$$

$$G_{3t} = G_3^2 x, \quad G_{3xx} = -G_3, \quad G_{1x} = G_1 x.$$

One can find the particular solution for equations in (41) as

$$G_1 = e^{\frac{x^2}{2}}, \quad G_3 = 0, \quad G_2 = 0, \quad K = e^{\frac{x^2}{2}}, \quad F = -t.$$

So that, one obtains the linearizing transformation

$$X = -t, \quad dT = e^{\frac{x^2}{2}} x' dt. \quad (42)$$

Hence, equation (40) is mapped by the transformation (42) into the linear equation

$$X'' = 0. \quad (43)$$

The general solution of equation (43) is

$$X = c_1 T + c_2, \quad (44)$$

where c_1 and c_2 are arbitrary constants. Applying the generalized linearizing transformation (42) to equation (44), we obtain that the general solution of equation (40) is

$$-t = c_1 \phi(t) + c_2,$$

where the function $T = \phi(t)$ is a solution of the equation

$$\frac{dT}{dt} = e^{\frac{x^2}{2}} x'.$$

D. The One-Dimensional Non-Polynomial Oscillator

In the note [13], Mathew and Lakshmanan presented a remarkable nonlinear system that all its bounded periodic motions are simple harmonic. The system is a particle obeying the highly nonlinear equation of motion

$$(1 + \lambda x^2)x'' + (\alpha - \lambda x'^2)x = 0, \quad (45)$$

where λ and α are arbitrary parameters.

By using our obtained theorems, we get the results as follow. Equation (45) is an equation of the form (8) in Theorem 2.1 with the coefficients

$$A_3 = 0, \quad A_2 = -\frac{\lambda x}{(\lambda x^2 + 1)}, \quad A_1 = 0, \quad A_0 = \frac{\alpha x}{(\lambda x^2 + 1)},$$

$$\lambda_1 = 0, \quad \lambda_2 = \alpha \lambda x(-\lambda x^2 + 2).$$

One can check that the condition (23) in Theorem 2.2. case (a) are satisfied. Now, the condition (26) is satisfied when the following condition holds, that is,

$$\alpha \lambda x(-\lambda x^2 + 2) = 0.$$

Two cases arise, that are $\alpha = 0$ and $\lambda = \frac{2}{x^2}$. (Note that for $\lambda = 0$ equation (45) is linear equation.)

Here we consider only case $\alpha = 0$. In this case, the equation (45) takes the form

$$(1 + \lambda x^2)x'' \lambda x x'^2 = 0. \quad (46)$$

The linearizing transformation is found by solving equations in Theorem 2.3 case (a), which become

$$F_t = -\frac{K}{G_1}, \quad K_x = -\frac{\lambda x K}{(1 + \lambda x^2)},$$

$$K_t = \frac{K(2G_{1t} \lambda x^2 + 2G_{1t} - \lambda x G_1 G_3)}{G_1(1 + \lambda x^2)}, \quad (47)$$

$$G_{3t} = -\frac{\lambda x^2 G_3^2}{(1 + \lambda x^2)}, \quad G_{3xx} = \frac{\lambda G_3(-\lambda x^2 + 1)}{(1 + \lambda x^2)^2},$$

$$G_{1x} = -\frac{\lambda x G_1}{(1 + \lambda x^2)}.$$

One can find the particular solution for equations in (47) as

$$G_1 = \frac{1}{(1 + \lambda x^2)^{\frac{1}{2}}}, \quad G_3 = 0, \quad G_2 = 0,$$

$$K = \frac{1}{(1 + \lambda x^2)^{\frac{1}{2}}}, \quad F = -t.$$

So that, one obtains the linearizing transformation

$$X = -t, \quad dT = \frac{1}{(1 + \lambda x^2)^{\frac{1}{2}}} x' dt. \quad (48)$$

Hence, equation (46) is mapped by the transformation (48) into the linear equation

$$X'' = 0. \quad (49)$$

The general solution of equation (49) is

$$X = c_1 T + c_2, \quad (50)$$

where c_1 and c_2 are arbitrary constants. Applying the generalized linearizing transformation (48) to equation (50), we obtain that the general solution of equation (46) is

$$-t = c_1 \phi(t) + c_2,$$

where the function $T = \phi(t)$ is a solution of the equation

$$\frac{dT}{dt} = \frac{1}{(1 + \lambda x^2)^{\frac{1}{2}}} x'.$$

E. Equation That Can Be Linearizable by Point and Sundman Transformations

Consider the nonlinear second-order ordinary differential equation

$$x'' + \mu_3 x^{k_3} x'^2 + \mu_2 x^{k_2} x' + \mu_1 x^{k_1} = 0, \quad (51)$$

where $k_3, k_2, k_1, \mu_1, \mu_2$ and $\mu_3 \neq 0$ are arbitrary constants. The Lie criteria [1], showed that the nonlinear equation (51) is linearizable by a point transformation if and only if $\mu_1 = 0$ and $\mu_2 = 0$. In [6], Nakpim and Meleshko showed that the nonlinear equation (51) is linearizable by a generalized Sundman transformation if and only if $\mu_2 \neq 0$ and $\mu_1 = 0$.

By using our obtained theorems, we get the results as follow. Equation (51) is an equation of the form (8) in Theorem 2.1 with the coefficients

$$A_3 = 0, A_2 = \mu_3 x^{k_3}, A_1 = \mu_2 x^{k_2}, \\ A_0 = \mu_1 x^{k_1}, \lambda_1 = k_2 \mu_2 x^{k_2},$$

$$\lambda_2 = x^{(k_1+k_3)} \mu_1 \mu_3 k_1 x + x^{(k_1+k_3)} \mu_1 \mu_3 k_3 x \\ + x^{k_1} \mu_1 k_1^2 + x^{k_1} \mu_1 k_1 - x^{2k_2} \mu_2^2 k_2 x.$$

Now, the conditions in Theorem 2.2. case (a) is satisfied when the following conditions holds, that are,

$$k_2 \mu_2 x^{k_2} = 0, \\ x^{(k_1+k_3)} \mu_1 \mu_3 k_1 x + x^{(k_1+k_3)} \mu_1 \mu_3 k_3 x + x^{k_1} \mu_1 k_1^2 \\ + x^{k_1} \mu_1 k_1 - x^{2k_2} \mu_2^2 k_2 x = 0.$$

Two cases arise.

Case 1: $\mu_2 = 0$ and $\mu_1 = 0$

In this case, the equation (51) takes the form

$$x'' + \mu_3 x^{k_3} x'^2 = 0. \quad (52)$$

The linearizing transformation is found by solving equations in Theorem 2.3 case (a), which become

$$F_t = -\frac{K}{G_1}, K_x = \mu_3 x^{k_3} K, \\ K_t = \frac{K(2G_{1t} + \mu_3 x^{k_3} G_1 G_3)}{G_1}, G_{3t} = \mu_3 x^{k_3} G_3^2, \\ G_{3xx} = -\frac{\mu_3 k_3 x^{k_3} G_3}{x}, G_{1x} = \mu_3 x^{k_3} G_1. \quad (53)$$

One can find the particular solution for equations in (53) as

$$G_1 = e^{\frac{\mu_3 x^{k_3+1}}{k_3+1}}, G_3 = 0, G_2 = 0, \\ K = e^{\frac{\mu_3 x^{k_3}}{k_3+1}}, F = -t.$$

So that, one obtains the linearizing transformation

$$X = -t, dT = e^{\frac{\mu_3 x^{k_3+1}}{k_3+1}} x' dt. \quad (54)$$

Hence, equation (52) is mapped by the transformation (54) into the linear equation

$$X'' = 0. \quad (55)$$

The general solution of equation (55) is

$$X = c_1 T + c_2, \quad (56)$$

where c_1 and c_2 are arbitrary constants. Applying the generalized linearizing transformation (54) to equation

(56), we obtain that the general solution of equation (52) is

$$-t = c_1 \phi(t) + c_2,$$

where the function $T = \phi(t)$ is a solution of the equation

$$\frac{dT}{dt} = e^{\frac{\mu_3 x^{k_3+1}}{k_3+1}} x',$$

where $k_3 \neq -1$.

For $k_3 = -1$, one can find the particular solution for equations in (53) as

$$G_1 = x^{\mu_3}, G_3 = 0, G_2 = 0, K = x^{\mu_3}, F = -t.$$

So that, one obtains the linearizing transformation

$$X = -t, dT = x^{\mu_3} x' dt. \quad (57)$$

Hence, equation (51) is mapped by the transformation (57) into the linear equation

$$X'' = 0. \quad (58)$$

The general solution of equation (58) is

$$X = c_1 T + c_2, \quad (59)$$

where c_1 and c_2 are arbitrary constants. Applying the generalized linearizing transformation (57) to equation (59), we obtain that the general solution of equation (51) is

$$-t = c_1 \phi(t) + c_2,$$

where the function $T = \phi(t)$ is a solution of the equation

$$\frac{dT}{dt} = x^{\mu_3} x'.$$

Case 2: $k_2 = 0$ and $\mu_1 = 0$

In this case, the equation (51) takes the form

$$x'' + \mu_3 x^{k_3} x'^2 + \mu_2 x' = 0. \quad (60)$$

The linearizing transformation is found by solving equations in Theorem 2.3 case (a), which become

$$F_t = -\frac{K}{G_1}, K_x = K \mu_3 x^{k_3}, \\ K_t = \frac{K(2G_{1t} + \mu_3 x^{k_3} G_1 G_3 - \mu_2 G_1)}{G_1}, \\ G_{3t} = G_3(x^{k_3} G_3 \mu_3 - \mu_2), G_{3xx} = -\frac{G_3 \mu_3 k_3 x^{k_3}}{x}, \\ G_{1x} = G_1 \mu_3 x^{k_3}. \quad (61)$$

One can find the particular solution for equations in (61) as

$$G_1 = e^{\frac{\mu_3 x^{k_3+1}}{k_3+1}}, G_3 = 0, G_2 = 0, \\ K = e^{\frac{\mu_3 x^{k_3}}{k_3+1} - \mu_2 t}, F = \frac{e^{\mu_2 t}}{\mu_2}.$$

So that, one obtains the linearizing transformation

$$X = \frac{e^{\mu_2 t}}{\mu_2}, dT = e^{\frac{\mu_3 x^{k_3+1}}{k_3+1}} x' dt. \quad (62)$$

Hence, equation (60) is mapped by the transformation (62) into the linear equation

$$X'' = 0. \quad (63)$$

The general solution of equation (63) is

$$X = c_1 T + c_2, \quad (64)$$

where c_1 and c_2 are arbitrary constants. Applying the generalized linearizing transformation (62) to equation (64), we obtain that the general solution of equation (60) is

$$\frac{e^{\mu_2 t}}{\mu_2} = c_1 \phi(t) + c_2,$$

where the function $T = \phi(t)$ is a solution of the equation

$$\frac{dT}{dt} = e^{\frac{\mu_3 x^{k_3+1}}{k_3+1}} x',$$

where $k_3 \neq -1$.

For $k_3 = -1$, one can find the particular solution for equations in (61) as

$$G_1 = x^{\mu_3}, \quad G_3 = 0, \quad G_2 = 0, \\ K = x^{\mu_3} e^{-\mu_2 t}, \quad F = \frac{e^{\mu_2 t}}{\mu_2}.$$

So that, one obtains the linearizing transformation

$$X = \frac{e^{\mu_2 t}}{\mu_2}, \quad dT = x^{\mu_3} x' dt. \quad (65)$$

Hence, equation (60) is mapped by the transformation (65) into the linear equation

$$X'' = 0. \quad (66)$$

The general solution of equation (66) is

$$X = c_1 T + c_2, \quad (67)$$

where c_1 and c_2 are arbitrary constants. Applying the generalized linearizing transformation (65) to equation (67), we obtain that the general solution of equation (60) is

$$\frac{e^{\mu_2 t}}{\mu_2} = c_1 \phi(t) + c_2,$$

where the function $T = \phi(t)$ is a solution of the equation

$$\frac{dT}{dt} = x^{\mu_3} x',$$

Remark 3.1: The conditions in Theorem 2.2. case (b) are satisfied if only if $\mu_1 = 0$.

F. Modified Generalized Vakhnenko Equation

In 2009, Ma, Li and Wang [14] focus on a modified generalized Vakhnenko equation (mGVE),

$$\frac{\partial}{\partial x} (L^2 u + \frac{1}{2} p u^2 + \beta u) + q L u = 0, \quad L = \frac{\partial}{\partial t} + u \frac{\partial}{\partial x}, \quad (68)$$

where ρ, q, β are arbitrary non-zero constants.

To develop the specific solutions for mGVE is exceedingly significant. For models, when $\rho = \beta = 0$ and $q = 1$, equation (68) is reduced to notable Vakhnenko equation (VE), which oversees the nonlinear engendering of high-recurrence wave in a loosening up medium [15]-[17]. The VE has soliton solutions [17]. When $\rho = q = 1$ and β an arbitrary non-zero constant, equation (68) is become as the generalized VE (GVE), in [18] it was indicated that GVE has N-soliton solution. When $\rho = 2q$ and β is an arbitrary non-zero constant, equation (68) has a loop-like, hump-like and cusp-like soliton solutions [19]. In [20], it was appeared that equation (68) has travelling wave solution and single-soliton solution.

Consider a modified generalized Vakhnenko equation (68), we can rewrite it in the form

$$2u_t u_{tx} + 2[uu_x u_{tx} + u_t (uu_{xx} + u_x^2)] + 2u^2 u_{xx} + 2uu_x^3 + \rho uu_x + \beta u_x + q(u_t + uu_x) = 0. \quad (69)$$

Of particular interest among solutions of equation (69) are travelling wave solutions:

$$u(t, x) = H(x - Dt),$$

where D is a constant phase velocity and the argument $x - Dt$ is a phase of the wave. Substituting the representation of a solution into equation (69), one finds

$$2D^2 H' H'' - 2DH'(2HH'' + H'^2) + 2H^2 H' H'' + 2HH'^3 + \rho H H' + \beta H' + q(-DH' + H H') = 0. \quad (70)$$

By using the obtained theorems, we get the results as follow. Equation (70) is an equation of the form in Theorem 2.1 with the coefficients

$$A_3 = 0, \quad A_2 = -\frac{1}{(D-H)}, \quad A_1 = 0, \\ A_0 = \frac{\rho H + \beta - qD + qH}{2(D^2 - 2DH + H^2)}, \quad \lambda_1 = 0, \quad \lambda_2 = \rho D + \beta.$$

From Theorem 2.2. case (a), equation (70) is linearizable if only if $\rho D + \beta = 0$.

G. Burgers' Equation

Burgers' equation is acquired because of the relationship between nonlinear wave movement and linear diffusion. It is the model for the investigation of consolidated impact of nonlinear advection and diffusion. The presence of the viscous term covers the wave-breaking, smooth out stun discontinuities, and thus we wish to get a tide and smooth solution. Also, as the dispersion term turns out to be vanishingly small, the smooth viscous solutions converge non-uniformly to the appropriate discontinuous shock wave, causing to another system for examining traditionalist nonlinear dynamical processes.

Consider the nonlinear convection-diffusion equation

$$\frac{\partial u}{\partial t} + u \frac{\partial u}{\partial x} - v \frac{\partial^2 u}{\partial x^2} = 0, \quad v > 0, \quad (71)$$

which is known as Burgers' equation. This equation balances between time advancement, nonlinearity, and dissemination. This is the nonlinear model equation for diffusive waves in fluid dynamics. Burgers (1948) first built up this equation basically to illuminate disturbance depicted by the collaboration of two inverse impacts of convection and dissemination.

The term uu_x will have a stunning up impact that will make waves break and the term vu_{xx} is a diffusion term like the one appearing in the heat equation.

Of particular interest among solutions of equation (71) are travelling wave solutions:

$$u(t, x) = H(x - Dt),$$

where D is a constant phase velocity and the argument $x - Dt$ is a phase of the wave. Substituting the representation of a solution into equation (71), one finds

$$-DH' + H H' - v H'' = 0. \quad (72)$$

By using the obtained theorems, we get the results as follow. Equation (72) is an equation of the form in Theorem 2.1 with the coefficients

$$A_3 = 0, A_2 = 0, A_1 = \frac{D-H}{v}, A_0 = 0, \\ \lambda_1 = \frac{1}{v}, \lambda_2 = \frac{-D+H}{v^2}.$$

One can check that these coefficients obey the condition in Theorem 2.2. case (b). Thus, equation (72) is linearizable via a generalized linearizing transformation.

IV. Conclusion

In this paper, the necessary condition which guarantee that the second-order ordinary differential equation can be linearized by generalized linearizing transformation is found in Theorem 2.1. Theorem 2.2 case (a) and case (b) are sufficient conditions for the linearization problem, they are selected by the value of λ_1 . A new algorithm for finding linearizing transformation is summarized in Theorem 2.3. Finally, some applications are provided to demonstrate our procedure.

References

- [1] S. Lie, "Klassifikation und Integration von gewöhnlichen Differentialgleichungen zwischen x, y , die eine Gruppe von Transformationen gestatten. III," *Archiv for Matematik og Naturvidenskab*, 8(4), 371-427, 1883.
- [2] R. Liouville, *Sur les Invariantes de Certaines Equations*, J. de l'École Polytechnique, 59, 7-88, 1889
- [3] A. M. Tresse, "Détermination des Invariants Ponctuels de l'Équation Différentielle Ordinaire du Second Ordre $y'' = f(x, y, y')$," *Preisschriften der fürstlichen Jablonowski schen Gesellschaft XXXII*, Leipzig, S.Herzel, 1896.
- [4] E. Cartan, *Sur les varietes a connexion projective*, Bull. Soc. Math. France, 52, 205-241, 1924.
- [5] L. G. S. Durate, I. C. Moreira and F. C. Santos, "Linearisation under nonpoint transformations," *J Phys A : Math Gen*, 27, L739-L743, 1994.
- [6] W. Nakpim and S. V. Meleshko, "Linearization of second-order ordinary differential equations by generalized Sundman transformations," *SIGMA*, 6, 051, 11 pages, 2010.
- [7] V. K. Chandrasekar, M. Senthilvelan and M. Lakshmanan, "A unification in the theory of linearization of second-order nonlinear ordinary differential equations," *Journal of Physics A: Mathematical and General*, 39, L69-L76, 2006.
- [8] V. K. Chandrasekar and M. Lakshmanan, "On the complete integrability and linearization of nonlinear ordinary differential equations," *Proc. R. Soc. A* 461, 2451-2477, 2005.
- [9] V. K. Chandrasekar and M. Lakshmanan, "On the complete integrability and linearization of nonlinear ordinary differential equations. II. Third-order equations," *Proc. R. Soc. A* 462, 1831-1852, 2006.
- [10] S. J. Farlow, *An introduction to differential equations and their applications*, Dover Publications, Inc., Newyork, 2012.
- [11] L. H. Koudahoun, J. Akande, D. K. K. Adjaï, Y. J. F. Kpomahou and M. D. Monsia, "Exact Classical and Quantum Mechanics of a Generalized Singular Equation of Quadratic Liénard Type," *Journal of Mathematics and Statistics*, Volume 14: 183.192, 2018.
- [12] M. T. Mustafa, A. Y. Al-Dweik and R. A. Mara'beh, "On the linearization of second-order ordinary differential equations to the laguerre form via generalized Sundman transformations," *symmetry, Integrability and Geometry: Methods and Applications (SIGMA)*, vol. 9, article no. 041, 2013.
- [13] P. M. Mathews and M. Lakshmanan, "On a unique nonlinear oscillator," *Q. Appl. Math.*, 32, 215-218, 1974.
- [14] Y. Ma, B. Li and C. Wang, "A series of abundant exact travelling wave solutions for a modified generalized Vakhnenko equation using auxiliary equation method," *Applied Mathematics and Computation*, 211, 102-107, 2009.
- [15] V.O. Vakhnenko and E.J. Parkes, "The two loop soliton solution of the Vakhnenko equation," *Nonlinearity*, 11, 1457-1464, 1998.
- [16] V.O. Vakhnenko, "High-frequency soliton-like waves in a relaxing medium," *J. Math. Phys.*, 40, 2011-2020, 1999.
- [17] V.O. Vakhnenko, E.J. Parkes, "The calculation of multi-soliton solutions of the Vakhnenko equation by the inverse scattering method," *Chaos, Solitons and Fractals*, 13, 1819-1826, 2002.
- [18] A.J. Morrison, E.J. Parkes, "The N-soliton solution of the modified generalized Vakhnenko equation (a new nonlinear evolution equation)," *Chaos, Solitons and Fractals*, 16, 13-26, 2003.
- [19] V.O. Vakhnenko, E.J. Parkes, "A Bäcklund transformation and the inverse scattering transform method for the generalised Vakhnenko equation," *Chaos, Solitons and Fractals*, 17, 683-692, 2003.
- [20] Y.P. Liu, Z.B. Li, K.C. Wang, "Symbolic computation of exact solutions for a nonlinear evolution equation," *Chaos, Solitons and Fractals*, 31, 1173-1180, 2007.

Prakrong Voraka received her B.S. degree in mathematics from Khon Kaen University, Khon Kaen, Thailand, in 2001 and her M.S. degree in mathematics from Kasetsart University, Bangkok, Thailand, in 2003. She received Ph.D. degree in applied mathematics in 2008 from Suranaree University of Technology, Nakhonratchasima, Thailand. Presently, she is a professor with the Department of General Science, Faculty of Science and Engineering, Kasetsart University Chalermphrakiat Sakon Nakhon Province Campus, Sakon Nakhon, Thailand. Her research interests are Lie group analysis, linearization problems and differential equations.

Supaporn Suksern received her B.S. degree in mathematics from Chiangmai University, Chiangmai, Thailand, in 2001 and her M.S. degree in applied mathematics from Chiangmai University, Chiangmai, Thailand, in 2004. She received Ph.D. degree in applied mathematics in 2009 from Suranaree University of Technology, Nakhonratchasima, Thailand. Presently, she is a professor with the Department of Mathematics, Faculty of Science, Naresuan University, Phitsanulok, Thailand. Her research interests are linearization problems and differential equations.

Nontakan Donjiwprai received his B.S. degree in mathematics from Naresuan University, Phitsanulok, Thailand, in 2020.

ตารางเปรียบเทียบกิจกรรมที่วางแผนไว้ และกิจกรรมที่ดำเนินการมา
กิจกรรมที่วางแผนไว้

กิจกรรม	เดือนที่												
	1	2	3	4	5	6	7	8	9	10	11	12	
1. ศึกษาโครงสร้างของสมการเชิงอนุพันธ์สามัญ อันดับสอง การแปลงในรูปแบบต่างๆ และ ผลงานวิจัยที่เกี่ยวข้องที่มีนักวิจัยทำมาก่อนหน้า นี้	↔												
2. ค้นคว้าหาเอกสาร ตำรา วารสาร และ เอกสารสิ่งพิมพ์ที่เกี่ยวข้องกับงานวิจัยที่กำลัง ดำเนินการวิจัยอยู่จากแหล่งข้อมูลต่างๆ		↔											
3. คิดค้นและวิจัยเพื่อหาเงื่อนไขที่จำเป็นสำหรับ การทำให้เป็นเชิงเส้น			↔										
4. คิดค้นและวิจัยเพื่อหาเงื่อนไขที่เพียงพอ สำหรับการทำให้เป็นเชิงเส้น				↔									
5. คิดค้นและวิจัยเพื่อหาการแปลงเชิงเส้น						↔							
6. สร้างโปรแกรมสำเร็จรูปในการทดสอบความ เป็นเชิงเส้น							↔						
7. คิดค้นและวิจัยเพื่อหาตัวอย่างและการ ประยุกต์ใช้								↔					
8. เขียนและพิมพ์ผลงานวิจัยเพื่อส่งพิจารณา ตีพิมพ์									↔				
9. รายงานสรุปผลโครงการ												↔	

ผลที่ได้รับตลอดโครงการ

ได้ตีพิมพ์ผลงานในวารสารวิชาการระดับนานาชาติในฐานข้อมูลของ SJR อยู่ใน Quartile 3 จำนวน 1 เรื่อง ได้แก่

Prakrong Voraka, Supaporn Suksern, and Nontakan Donjiwprai. Linearizability of Nonlinear Second-Order Ordinary Differential Equations by Using a Generalized Linearizing Transformation. IAENG International Journal of Applied Mathematics., 50:4, IJAM_50_4_19

