




MORPHOLOGICAL AND RIBOSOMAL DNA- BASED CHARACTERIZATION  
OF COMMERCIAL *CORDYCEPS MILITARIS* STRAINS IN THAILAND




JUNYA NOPPARAT

A Thesis Submitted to the Graduate School of Naresuan University  
in Partial Fulfillment of the Requirements  
for the Master of Science in (Agricultural Biotechnology)

2018

Copyright by Naresuan University

MORPHOLOGICAL AND RIBOSOMAL DNA- BASED CHARACTERIZATION  
OF COMMERCIAL *CORDYCEPS MILITARIS* STRAINS IN THAILAND



A Thesis Submitted to the Graduate School of Naresuan University  
in Partial Fulfillment of the Requirements  
for the Master of Science in (Agricultural Biotechnology)  
2018

Copyright by Naresuan University

Thesis entitled "Morphological and ribosomal DNA- based characterization of commercial *Cordyceps militaris* strains in Thailand"

By JUNYA NOPPARAT

has been approved by the Graduate School as partial fulfillment of the requirements  
for the Master of Science in Agricultural Biotechnology of Naresuan University

**Oral Defense Committee**

..... Chair  
(Associate Professor Touchkanin Jongjitvimol, Ph.D.)

..... Advisor  
(Assistant Professor Wassana Chatdumrong, Ph.D.)

..... Co Advisor  
(Assistant Professor Kawee Sujipuli, Ph.D.)

..... Co Advisor  
(Assistant Professor Surisakk Prasarnpun, Ph.D.)

..... Internal Examiner  
(Dr. Thanita Boonsrangsom)

**Approved**

.....  
(Professor Paisarn Muneesawang, Ph.D.)

for Dean of the Graduate School

<b>Title</b>	MORPHOLOGICAL AND RIBOSOMAL DNA-BASED CHARACTERIZATION OF COMMERCIAL <i>CORDYCEPS MILITARIS</i> STRAINS IN THAILAND
<b>Author</b>	JUNYA NOPPARAT
<b>Advisor</b>	Assistant Professor Wassana Chatdumrong, Ph.D.
<b>Co-Advisor</b>	Assistant Professor Kawee Sujipuli, Ph.D. , Assistant Professor Surisakk Prasarnpun, Ph.D.
<b>Academic Paper</b>	Thesis M.S. in Agricultural Biotechnology, Naresuan University, 2018
<b>Keywords</b>	<i>Cordyceps militaris</i> , Morphology, Ribosomal DNA

### ABSTRACT

*Cordyceps militaris*, an entomopathogenic fungus, has been extensively utilized for many years as functional food and for medicinal remedies in Thailand. The fungus has numerous strains which are commonly sold in herbal markets and not all have been identified. Seven commercial *C. militaris* isolates (*Cmi1*, *Cmi2*, *Cmi3*, *Cmi4*, *Cmi5*, *Cmi6* and *Cmi7*) were studied using macroscopic studies. The stromatal colour of seven isolates were golden yellow. Of these, six isolates had a cylindrical stipe of stromata (approximately 0.3-0.4 cm wide x 4-6 cm long) with a round cap, except the *Cmi4*, stroma had the widest stipe (approximately 1.5-2.5 cm wide and 2-4 cm long) with a shape cap. Moreover, the mycelium had the highest mycelial growth rate, but only the *Cmi4* had the lowest. To determine the stromatal characteristics using cross section technique and observing under a microscope, the result revealed that all isolates were golden yellow at the edge, and colourless at hyphae. To investigate the spore characteristics using the spore shooting technique, and observing under a microscope, the result revealed that most spores were globular shape. To study the conidial formations and hyphae characteristics using the slide culture and SEM techniques, the result showed that most spores had globose and oval shapes, moreover these germinated and produced long hyphae. The conidia formed mostly single globe heads and rare group globe heads in short chains and subtending conidia. Some conidia had pyriform or cylindrical shapes. Genetic variation among

the seven isolates gave 100% nucleotide-sequence homology at the ITS and 28S rDNA regions. These data suggested that the macroscopic method was a highly powerful tool for strain identification of *C. militaris*.



## ACKNOWLEDGEMENTS

I would like to offer my profound gratitude to my advisor Assistant Professor Dr. Wassana Chatdumrong who had being tirelessly coaching me through this work and co-advisor Assistant Professor Dr. Kawee Sujipuli and Assistant Professor Dr. Surisak Prasarnpun for constant support during the studies.

I would like also to give special thanks to all the thesis committee members and appreciate deeply their constructive comments that shape this writing.

I would like to acknowledge the Agricultural Research Development Agency (ARDA; Project number CRP5805020130).

I would like also to appreciate the Department of Agricultural Science and Center of Excellence in Research for Agriculture Biotechnology (CERAB), Faculty of Agriculture, Natural Resources and Environment, Department of Microbiology and Parasitology, Faculty of Medical Science, and The Department of Biology, Faculty of Science for support laboratory instruments and facility.

I would like to acknowledge every staff for their guidance especially Assistant Professor Dr. Anupan Kongbangkerd, Mr. Boworn Kunakhonnurak, Mr. Sompop Pinit, Miss Kamria Woranoot, Mr. Teerawech Promchiangsa and Miss Chutiporn Maneeply.

I would like to register very special thanks to my father (Mr. Phongsakorn Nopparat), my mother (Mrs Pat Nopparat), my sister (Miss Chutima Nopparat) and my lovely boyfriend (Mr. Nigel Roberto John Smith) for their supports.

Finally, I would like to thanks all of my colleagues and senior graduate students and all those who are not mentioned here for their support in different capacities for the success of this thesis.

JUNYA NOPPARAT

## TABLE OF CONTENTS

	Page
ABSTRACT .....	C
ACKNOWLEDGEMENTS .....	E
TABLE OF CONTENTS .....	F
CHAPTER I INTRODUCTION .....	1
1.1 General introduction .....	1
1.2 Objectives of the study .....	2
1.3 Research Significance .....	2
1.4 Research Scope .....	2
1.5 Keywords .....	3
1.6 Research Hypothesis .....	3
CHAPTER II LITERATURE REVIEW .....	4
2.1 Kingdom fungi .....	4
1) Phylum Chytridiomycota .....	5
2) Phylum Zygomycota .....	5
3) Phylum Basidiomycota .....	5
4) Phylum Ascomycota .....	5
2.2 Phylum Ascomycota .....	5
2.3 <i>Cordyceps</i> group .....	7
2.4 Source of <i>Cordyceps</i> .....	8
2.5 <i>C. militaris</i> .....	9
2.6 Benefits of <i>C. militaris</i> .....	11
2.8 Effect of parameters on <i>C. militaris</i> cultivation .....	12
2.9 Morphological characteristics study of <i>Cordyceps</i> .....	13
2.10 Molecular markers used for genotyping fungi .....	15
2.9 Definition of Internal transcribed spacer (ITS) and 28S ribosomal DNA .....	15




2.10 Identification of <i>Cordyceps</i> using ITS and 28S rDNA markers.....	16
CHAPTER III MATERIALS AND METHODS .....	19
3.1 Macroscopic studies.....	19
1) Macroscopic characters of <i>C. militaris</i> .....	19
2) Mycelium growth.....	19
1) Cross section of stromata.....	19
2) Spore shooting.....	20
3) Slide culture technique .....	20
4) SEM Study .....	20
3.3 Genomic DNA (gDNA) extraction and PCR amplification .....	21
3.4 DNA sequence alignment .....	21
CHAPTER IV RESULTS AND DISCUSSION.....	23
4.1 Macroscopic studies.....	23
1) Macroscopic characters of <i>C. militaris</i> .....	23
2) Mycelium growth.....	24
4.2 Microscopic characteristics studies .....	27
1) Microscopic characteristics of stromata among <i>C. militaris</i> .....	27
2) Cross section of stromata among <i>C. militaris</i> .....	28
3) Spore shooting technique.....	29
4) Slide culture technique .....	30
5) Observation of conidial formation by Scanning Electron Microscopy (SEM) .....	31
4.3 DNA sequencing.....	32
CHAPTER V CONCLUSION .....	38
Suggestion.....	39
REFERENCES.....	40
APPENDIX .....	48
BIOGRAPHY .....	59



## LIST OF TABLES

<b>Table 1</b> Host species of <i>C. militaris</i> belonging to different order and families of insect.....	10
<b>Table 2</b> Primers used to amplify fungal 28S rDNA genes and ITS region sequences	22
<b>Table 3</b> Mycelial growth characteristics of seven <i>C. militaris</i> isolates ( <i>Cmi1-Cmi7</i> ) on PDA .....	26
<b>Table 4</b> Colony diameter of seven <i>C. militaris</i> ( <i>Cmi1- Cmi7</i> ) isolates on PDA medium .....	26
<b>Table 5</b> Colony pigmentation of <i>C. militaris</i> isolates ( <i>Cmi1- Cmi7</i> ) on PDA medium	27
<b>Table 6</b> Blast results of the ITS nucleotide sequences generated from studied .....	35
<b>Table 7</b> Blast results of the 28S rDNA nucleotide sequences generated from studied	37



## LIST OF FIGURES

<b>Figure 1</b> Two types of hyphae characteristics, septate hyphae (left) and non-septate hyphae (right) .....	4
<b>Figure 2</b> Life cycle of fungi in phylum Ascomycota.....	6
<b>Figure 3</b> Life cycle of <i>Cordyceps</i> .....	7
<b>Figure 4</b> Morphology of <i>Cordyceps</i> .....	8
<b>Figure 5</b> Map positions of primer pairs (ITS1-ITS4 and LROR-LR7) used to amplify PCR products .....	16
<b>Figure 6</b> Morphological characteristic of stromata among <i>C. militaris</i> isolates ( <i>Cmi1-Cmi7</i> ) .....	24
<b>Figure 7</b> Mycelial growth rates and characteristics among seven <i>C. militaris</i> isolates ( <i>Cmi1</i> to <i>Cmi7</i> ) cultured on PDA medium for 3, 6, 9, 12, 15, 18 and 21 days.....	25
<b>Figure 8</b> The outer surface characteristics of stromata among seven <i>C. militaris</i> isolates observed under Stereo microscope .....	28
<b>Figure 9</b> Evaluation of stromal characteristics among seven <i>C. militaris</i> isolates ( <i>Cmi1</i> to <i>Cmi7</i> ) by cross section technique under a compound microscope .....	29
<b>Figure 10</b> Spore production of <i>C. militaris</i> <i>Cmi1-Cmi7</i> isolates by spore shooting ...	30
<b>Figure 11</b> Spore germination of different isolates of <i>C. militaris</i> <i>Cmi1-Cmi7</i> by slide culture technique for 14 days, (A) Conidia formed in short chains (B) Conidia formed in globose head.....	31
<b>Figure 12</b> Scanning electron micrographs of conidial hyphae of <i>C. militaris</i> <i>Cmi1-Cmi7</i> 14 days after culture on PDA (Scales: A = 1 $\mu$ m, B = 5 $\mu$ m).....	32
<b>Figure 13</b> Comparative analysis of nucleotide sequences between seven studied <i>C. militaris</i> isolates and <i>C. militaris</i> NBRC100741 strain using ClastaW program .....	34
<b>Figure 14</b> Comparative analysis of nucleotide sequences between seven studied <i>C. militaris</i> isolates and <i>C. militaris</i> JM0807 strain using ClastaW program.....	37

# CHAPTER I

## INTRODUCTION

### 1.1 General introduction

*Cordyceps militaris* is an entomopathogenic fungus whose spore invades and grows in a specific insect host in caterpillar stage which hibernates underground during the winter season. Later, during the summer season, the spore develops into a fruiting-body formation on the surface of the insect's cadaver (Guo et al., 2017). In Chinese traditional medicine, the stroma (fruiting body) of *C. militaris* is credited with several pharmacological properties, and widely used as a herb, functional food, for improvement of immunity (Izgi et al., 2015; J.-y. Liu et al., 2016; Song et al., 2016), for bacteriostasis (Avtonomova, Krasnopolskaya, Shuktueva, Isakova, & Bukhman, 2015; Chiu et al., 2016), antitumor (P. Zhang et al., 2015), antiasthmatic effects (Tianzhu, Shihai, & Juan, 2015), and as antihyperuricemic (Yong et al., 2016) and antihypoglycemic (L. Ma, Zhang, & Du, 2015). Main abundant bioactive compounds produced by various *Cordyceps* species were adenosine (Tatani et al., 2016), ergosterol, mannitol and exopolysaccharide (Raethong, Laoteng, & Vongsangnak, 2018), and cordycepin (Yong et al., 2016). Of these, cordycepin is an extremely valuable bioactive compound produced only by *C. militaris*. Artificial cultivation gives higher production of cordycepin compared to nature; however, its quality and quantity depend on fungal culture media (Lin, Lai, Wu, Hsu, & Tai, 2018), culture conditions (G.-H. Sung, Shrestha, Han, & Sung, 2011), and isolates (T. Wen, Long, Kang, Wang, & Zeng, 2017). Recently, *C. militaris* has been isolated and estimated to many isolates in Asian counties (Kang et al., 2017) including in Thailand (Tapingkae, 2012), where isolates contained various amounts of cordycepin (T. Wen et al., 2017). Therefore, classification among isolates of *C. militaris* was beneficial to screen for isolates that were able to produce high cordycepin content. During the past decade, two methods were widely used for morphological classification of *C. militaris* isolates consisting of macroscopic such as morphological characteristics (such as shape, size, and colour of stroma) (H.-j. Liu, Hu, Chu, Li, & Li, 2011) and mycelium

growth (Dang, Wang, & Lay, 2018). Addition, microscopic such as cross section (Guo et al., 2017), spore shooting technique (Choi, Hyde, & Ho, 1999), slide culture technique (Leck, 1999) and SEM techniques (Albini, 2017). Recently, molecular markers corresponding to internal transcribed spacer (ITS) (Lam et al., 2015) and 28S-ribosomal DNAs (rDNAs) regions (Y. Li, Jiao, & Yao, 2013) has gained prominence to classify among *Cordyceps* isolates. Therefore, the aim of this study was to identify Thai commercial isolates of *C. militaris* using both morphological characteristics and molecular markers. Results will be beneficial by providing the accurate method for identifying commercial isolates of Thai medicinal fungus *C. militaris*.

## 1.2 Objectives of the study

- 1) To study morphological characteristic of *C. militaris* isolates from Thai commercial sources.
- 2) To study nucleotide variation at rDNA regions using ITS- and 28S-markers.

## 1.3 Research Significance

The morphological characteristic and genetic variation among *C. militaris* isolates are highly powerful methods to identify among isolates of *C. militaris* in Thai commercial strains.

## 1.4 Research Scope

*C. militaris* were collected from seven different commercial farms in Thailand. The methods used for identifying and studying *C. militaris* isolates were morphological characteristics (such as shape, size, and colour of stroma), growth development on media, cross section of stroma, spore shooting technique, slide culture technique, and SEM technique. Genomic DNA was isolated using InstaGene Matrix (Bio-Rad, United States) kit, quantified using gel electrophoresis system and spectrophotometric system. ITS and 28S rDNA markers were used for DNA fingerprinting using PCR techniques. Purified DNA samples were performed using a PRISM BigDye Terminator v3.1 Cycle Sequencing Kit (Applied Biosystems, Foster

City, Canada) and then sequenced and analyzed using an ABI Prism 3730XL DNA analyzer (Applied Biosystems, Foster City, Canada).

### 1.5 Keywords

*Cordyceps militaris*, Morphology, Ribosomal DNA

### 1.6 Research Hypothesis

Macroscopic, microscopic and molecular marker methods would be proficient tools for morphological classification of *C. militaris* isolates in Thailand.




## CHAPTER II

### LITERATURE REVIEW

Fungi, eukaryotic organisms, have one or more nucleus enclosed within membrane without chlorophyll. Therefore, they cannot photosynthesize their cells. Heterotrophic fungi absorb any organic matters from external sources and environment. Moreover, they may exist as parasites obtaining nutrients from a living host. Fungal cell wall consists of cellulose, hemicellulose and chitin compound (Suwanpinit & Suwanpinit, 1998).

Generally, the structures of fungi are made up of thread like filaments and colourless hyphae. Its functions are associated with growth development, and reproduction by forming spores. The hyphal tip region is the highest growth rate, and many hyphae are formed to mycelium (5-10  $\mu\text{m}$  diameter) (Sangtian, 2009). Their cell structures of hyphae are different characteristics which can be divided into 2 types, septate hyphae and non-septate hyphae (Figure 1).



**Figure 1 Two types of hyphae characteristics, septate hyphae (left) and non-septate hyphae (right) (Alexopoulos, Mims, & Blackwell, 1996)**

#### 2.1 Kingdom fungi

Dube (2013) reported that the fungus belongs to the Kingdom Fungi which can be divided of various phylum (Sylvia S. Mader et al., 2007) as following:



### **1) Phylum Chytridiomycota**

Most fungi in this phylum have small thallus formations, producing sporangiums, and have root like structures, called rhizoids. The general characteristic of this fungus has a single flagellum, which is using for movement. In addition, the flagellum is able to differentiate to a zoospore with one nucleus, used as sexual-reproduction systems such as planogametic copulation, gametangial copulation and somatogamy.

### **2) Phylum Zygomycota**

The fungi in this phylum are often referred to lower fungi with having a single cell and non-septate. Most fungi can grow and live in the water, soil, and fossil (decaying plant or animal material). Some fungi are parasites to plants, insects, and small animals, while others are symbiotic relationships with plants. In an asexual reproductive stage, a hundreds of haploid spores are developed from a sporangium at the tip of upright hyphae.

### **3) Phylum Basidiomycota**

The fungi in this phylum are often referred to higher fungi with having septate hyphae, packing to a stick like a stem such as fruiting body. As an asexual reproductive stage, the conidia are formed at the conidiophore. As a sexual reproductive stage, the spore is formed at the basidium, called basidiospore.

### **4) Phylum Ascomycota**


Its members are commonly, known as the sac fungi or ascomycetes, and characterized by a saclike structure. The ascus contains four to eight ascospores in the sexual stage. For an asexual reproductive stage, the conidia are formed at the end of hyphae.

## **2.2 Phylum Ascomycota**

Phylum Ascomycota (sac fungi) is classified as higher fungus, consisting of approximately 60,000 species, which are extensively found in all land ecosystems. Their spores and hyphal fragments are found in the atmosphere, in fresh-water, and in salt-water environments. In this phylum, the largest class is ascomycetes, including



the morel (cup fungi) and the truffle (mushroom). Some fungi in class ascomycetes (in phylum Ascomycota) are plant pathogens, animal pathogens, edible mushrooms, and living on dead organic matter (as saprobes). The unique characteristics of ascomycetes are spore formation, which is in a sac-like structure called an ascus. During sexual reproduction, thousands of asci are located in a large stroma called the ascocarp (Figure 2). The ascus contains four to eight ascospores in the sexual stage. For an asexual reproductive stage, the conidia are formed at the end of hyphae. Mycelia are composed of septate hyphae. The hyphae of ascomycetes consists of homokaryotic hypha or heterokaryotic hypha and pseudomycelium (Sangtian, 2009). The economic importance of ascomycetes is used in food production, such as yeast, which is used to produce alcohol and make bread rise. Ascomycetes are used in medicine as well, as it can see with the antibiotic penicillin. Some ascomycete fungi exist as parasites with obtaining nutrients from a living host such as insects. Insect's fungi in this class are considered important to the economy. It has been a highly valued traditional medicines such as the genus of *Ophiocordyceps*, *Isaria* and especially *Cordyceps* (Tapingkae, 2012).




**Figure 2 Life cycle of fungi in phylum Ascomycota**

Note: This illustration shows fungal life cycle either asexual (left) or sexual (right) reproduction (Sangtian, 2009)

### 2.3 *Cordyceps* group

*Cordyceps* group is commonly referred to as Chinese caterpillar fungus or winter-worm and summer-grass, which is a literal translation from its Chinese name, Dong Chong Xia Cao. *Cordyceps* is an entomopathogenic fungus whose spore invades and grows in a specific insect host in a caterpillar stage which hibernates underground covered with snow during the winter season. Later, during the summer season, the spores grow and absorb nutrient and mineral from the insect host, and develops into a stroma formation on the surface of the insect's cadaver. The stroma has been developed to mushroom, shooting out many spores for sexual reproduction (Guo et al., 2017) (Figure 3). In nature, *Cordyceps* is consist of head of stroma, stalk (look like mushroom) and larva (Buenz, Bauer, Osmundson, & Motley, 2005; Shrestha, Zhang, Zhang, & Liu, 2010) (Figure 4).



**Figure 3 Life cycle of *Cordyceps***

Note: This illustration shows Life cycle of *Cordyceps* (Tapingkae, 2014)



**Figure 4 Morphology of *Cordyceps***

Note: This illustration shows morphology of *Cordyceps* in the nature (Tapingkae, 2014)

#### **2.4 Source of *Cordyceps***

C. Li et al. (2006) reported that *Cordyceps* were classified more than 750 species around the world. Out of these, the most species (approximately 400 species) have been found in Asia (Nepali, China, Japan, Korea, Vietnam and Thailand), because its majority regions are various humid climate regions and tropical forest (G.-H. Sung et al., 2007). In Thailand, Sivichai (2001) found more than 80 *Codyceps* species. Tapingkae (2012) studied diversity of *Codyceps* at Doi Suthep-Pui National Park in Chiang Mai, found that genus of *Codyceps* belongings to family Clavicipitaceae, consisting of many *Codyceps* species which were parasites in various insect varieties. The various hosts of *Cordyceps* consist of ant, bee, wasp, hornet, spider, aphids, grub, dragonfly, butterfly and worm etc. Generally, insect fungi are specific to particular host species (Sung et al., 2007). Insect fungi in genus of *Cordyceps* was used to pharmaceutical industry as food for humans and not allergic (G.-H. Sung et al., 2007). The important *Cordyceps* in this genera are *Ophiocordyceps sinensis*, *Isaria tenuipes*, *Isaria sinclairii* and *C. militaris* (Tapingkae, 2012).

### 2.5 *C. militaris*

*C. militaris* can be classified according to taxonomy as follows (Das, Masuda, Sakurai, & Sakakibara, 2010).

Kingdom	Fungi
Phylum	Ascomycota
Sub-phylum	Ascomycotina
Class	Ascomycetes/ Pyrenomycetes
Order	Hypocreales
Family	Clavicipataceae
Genus	<i>Cordyceps</i>
Species	<i>Cordyceps militaris</i>

*C. militaris*, as an entomopathogenic fungus, infects many insects, especially most Lepidoptera larvae of different host species as shown more detail in Table 1 (Tapingkae, 2012).

**Table 1** Host species of *C. militaris* belonging to different order and families of insects

Order	Family	Species	References
Coleoptera	Scarabaeidae	<i>Lachnosterna quercina</i>	(Farlow & Seymour, 1888)
	Scolytidae	<i>Ips sexdentatus</i>	(Kryukov, Yaroslavtseva, Lednev, & Borisov, 2011)
Diptera	Tenebrionidae	<i>Tenebrio molitor</i>	(De Bary, 1867)
	Tipulidae	<i>Tipula paludosa</i>	(Müller-Kögler, 1964)
Hymenoptera	Cimbicidae	<i>Cimbex similis</i>	(Y. Kobayasi, 1941)
Lepidoptera	Arctiidae	<i>Euprepia caja</i>	(De Bary, 1867)
	Bombycidae	<i>Andraca bipunctata</i>	(Panigrahi, 1995)
		<i>Bombyx caja</i>	(Cesati, 1861)
		<i>B. mori</i>	(Gu & Liang, 1987)
		<i>B. pythiocampa</i>	(Y. Kobayasi, 1941)
		<i>B. rubi</i>	(Tulasne, 1865)
	Drepanidae	<i>Cymatophora duplaris</i>	(Lagerberg, 1922)
		<i>C. flavicornis</i>	(Lagerberg, 1922)
		<i>Palimpsestis duplaris</i>	(Ulvinen, 1969)
		<i>Polyploca flavicornis</i>	(Ulvinen, 1969)
	Geometridae	<i>Biston 10alcate</i>	(T. Ma et al., 2007)
		<i>Culcula panterinaria</i>	(Qiwu, 1997)
		<i>Lycia hirtaria</i>	(Ulvinen, 1969)
	Hepialidae	<i>Hepialus sp.</i>	(Petch, 1948)
	Lasiocampidae	<i>Dendrolimus latipennis</i>	(Jing, 1987)
		<i>D. pini</i>	(Sopp, 1911)
		<i>D. sibiricus and Macrothylacia rubi</i>	(Kryukov et al., 2011)
		<i>Gastropacha quercus</i>	(De Bary, 1867)
	Lymantriidae	<i>Dasychira pudibunda</i>	(Ulvinen, 1969)
		<i>Leucoma salicis</i>	Kryukov et al. 2011
	Notodontidae	<i>Fentonia ocypete,</i>	(Qiwu, 1997)
		<i>Lampronadata cristata,</i>	
		<i>Phalera assimilis and Phalerodonta albibasis</i>	
		<i>Phalera bucephala</i>	
		<i>Quadricalcarifera punctatella</i>	(Gray, 1858)
		<i>punctatella</i>	(Sato, Shimazu, & Kamata, 1994)
		<i>punctatella</i>	



**Table 1 (Cont.)**

Order	Family	Species	References
	Saturniidae	<i>Anisota senatoria</i>	(Hitchcock, 1961)
	Sphingidae	<i>Callambulyx tatarinovi</i>	(Y. Kobayasi, 1941)
		<i>Laothë populi</i>	(Ulvinen, 1969)
		<i>Marumba sperchius</i>	(Qiwu, 1997)
		<i>Mimas tiliae</i>	(Ulvinen, 1969)
		<i>Sphinx euphorbiae</i>	(De Bary, 1867)
		<i>S. pinastri</i>	(Ulvinen, 1969)
	Thyatiridae	<i>Tetheella fluctuosa</i>	(Kryukov et al., 2011)
		<i>Ochropacha duplaris</i>	
		<i>Tethea ocularis</i>	

## 2.6 Benefits of *C. militaris*

Nowadays, the *Cordyceps* was publically used as safely herbal products, which has gained popular increase in the United States, Japan, Australia, China, Switzerland, Korea, Taiwan, Malaysia, Canada, Hong Kong and Thailand (Das et al., 2010). It had previous reported that the *C. militaris* and the *C. sinensis*, now known as *O. sinensis* (G.-H. Sung et al., 2007), had high Cordycepin, which was major bioactive compound with higher than other types of spices (Dong, Lei, Ai, & Wang, 2012). Minor bioactive compounds, found in *C. militaris*, consist of monosaccharide, disaccharide, polysaccharide, beta-glucan, mannitol, galactose, adenosine, cordycepic, amino acid and vitamin had been found in *C. militaris* (Dong et al., 2012), including organic matters of protein, lipid, vitamin, mineral and amino acid (Tapingkae, 2013). The *Cordyceps* had various mineral such as potassium, sodium, calcium, magnesium, iron, copper, manganese, zinc, phosphorus and selenium (Bhandari et al., 2010). The digested enzymes related in monogastric simple nonruminants *C. militaris* found in *C. militaris* are amylase, pectinase,  $\beta$ -glucanase, endoglucanase, cellulose, protease and phytase (Tapingkae, 2013). In Chinese traditional medicine, the stroma of *C. militaris* is credited with several pharmacological properties, and widely used as a herb, functional food, for improvement of immunity (Izgi et al., 2015; J.-y. Liu et al., 2016; Song et al., 2016), for bacteriostasis (Avtonomova et al., 2015; Chiu et al., 2016), antitumor (P. Zhang et al., 2015), antiasthmatic effects (Tianzhu et al., 2015), and as a antihyperuricemic (Yong et al., 2016) and antihypoglycemic (L. Ma et al., 2015).

## 2.8 Effect of parameters on *C. militaris* cultivation

The *C. militaris* cultivation in *in vitro* conditions were depending on various parameters such as culture medium, light-dark, temperature, pH, and humidity. Many previous studies have been reported as shown below:

Holliday, Cleaver, Loomis-Powers, and Patel (2004) have been developed the unique method for improving hybridized strains of *Cordyceps*, from wild collected *Cordyceps*, by using rattlesnake venom to enhance somatic fusion of two mycelial strains. These artificially hybridized strains were able to produce a higher and more consistent quality levels of adenocine and cordycepin compound.

Shrestha, Lee, Han, and Sung (2006) studied the growth of *C. militaris* on various media under light conditions. The result found that the light is most important for growth and pigmentation of mycelium. The hyphae had the highest growth under darkness condition, compared to light condition. Addition, Shrestha studied the effect of 22 culture media on growth of *C. militaris*. Of these, only three media which were Sabouraud Dextrose agar plus Yeast Extract (SDAY), Sabouraud Maltose agar plus Yeast Extract (SMAY) and Czapek Yeast Extract agar (CZYA) gave the highest growth development of *C. militaris*.

J.-M. Sung et al. (2006) and Pathania, Joshi, and Sagar (2015) reported that the suitable temperature at 25°C under darkness condition for 7 days gave the best mycelium growth of *C. militaris*.

Hong et al. (2010) reported that the appropriate condition for growing mycelial colonies and forming stroma of *C. militaris* was that mycelium infected on silkworm, incubated at 20°C under photoperiod at 1,000 lux and relative humidity 90% 9-10 days.

P.-D. Kang et al. (2010) referred to the popular method of culture *Isaria tennipes* in Korea using by silkworm (*Bombyx mori*). The larva of silkworm was feed in a tray in the evaporative cooling system. The starter of *Isaria tennipes* mixed with starch syrup and spray at 0, 12 and 24 hours on larva, which it was on 5 states of molting process (weakest moments). The larva was feed by leaf, and would go into pupa for 11 days. After that, the cocoon was cut and took the pupa out. The pupa (infected starter of *Isaria tennipes*) was incubated 20-22°C humidity 95% and would go into the stroma.



Kim, Shrestha, Sung, Han, and Sung (2010) studied optimum condition for regeneration of fruiting state and develop to the stroma of *Cordyceps*. The result showed that the optimum culture was incubated  $20 \pm 1^\circ\text{C}$  under the light condition and relative humidity 70- 80% for 60 days.

Pathania et al. (2015) studied the effect of pH in medium on mycelium growth of *Cordyceps* found that the mycelium had the highest growth on the medium at pH 7.

G.-H. Sung, Shrestha, Han, Kim, and Sung (2010) studied the effect of various carbon, nitrogen or mineral sources on mycelium growth of *Cordyceps*. The result showed carbon sources (fructose and xylose) gave the highest growth in comparison with other carbon sources (maltose, starch, dextrose, dextrin, sucrose, lactose and water agar), followed by nitrogen source (peptone) and mineral sources ( $\text{K}_2\text{HPO}_4$ ,  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$  and  $\text{NaCl}$ ). Similarly, G.-H. Sung et al. (2007) mention culture of *Cordyceps* genus can regenerate easily on liquid and solid medium supplement different concentration of carbon source and nitrogen source.

T. Wen, Li, Kang, Kang, and Hyde (2014) reported that various pH (5.0-8.0) conditions promoted stroma growth of *C. militaris*. The result revealed that the maximum stroma growth ( $1.81 \pm 0.08$  g/bottle) and the maximum cordycepin production ( $7.40 \pm 0.01$  mg/g dry weight) were accomplished at low pH ranged from 5.5 to 6.0, cultured on 40 g/L glucose, 5 g/L peptone, 1.5 g/L  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ , 1.5 g/L  $\text{K}_2\text{HPO}_4$  and 1.0 mg/L NAA for optimal stroma growth and 10 g/L glucose, 10 g/L peptone, 1.0 g/L  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ , 1.0 g/L  $\text{K}_2\text{HPO}_4$  and 1.0 mg/L NAA.

## 2.9 Morphological characteristics study of *Cordyceps*

The mycelium of anamorph stage, some can identify by observation of conidia formation on culture medium (asexual reproduction). Some cannot describe character of fungus through culture medium. Therefore, it is the reason that makes the morphology study quite difficult and has restriction. Generally, insect fungi in genus *Cordyceps* found sexual stage, which is period of time to produce stroma. Therefore, identification of this genus should be used the teleomorph state, which this state is produce ascospores inside of asci. The teleomorph state clearly to see a difference between species in *Cordyceps* such as shape, size, length and diameter of stroma. The

previous researches, since 1982-2013, have been successfully used to the identification of *Cordyceps* species using morphology techniques.

Y Kobayasi (1982) and Eriksson et al. (2003) identified *Cordyceps* species using morphological characteristics such as colour, shape of stroma and perithecia. The result found that these characteristics can be used to classify four species such as *C. subg. bola cordyceps*, *C. subg. cordyceps*, *C. subg. neocordyceps*, and *Cordyceps subg. Ophiocordyceps*

A. Liu (1997) and Shrestha, Han, Yoon, and Sung (2005) reported the approaches of different ascospore germination and conidial formation of *C. militaris* under the light microscope and electron microscope were utilized to determine *C. militaris* species. Generally, conidial formation was either solitary or whorl, subsequently the succeeding conidia was cylindrical and nearly globose which germinated to produced long hyphae.

Shrestha et al. (2005) studied the ascospore pigmentation of seven *C. militaris* strains infected on Lepidopteran pupae, collected from natural habitats. They were different colours such as yellow, orange and white, while the white colour (albino) was a mutant strain caused by gene mutation. The heritability of pigmentation characteristic is controlled by gene in parental strains, which this gene was transferred from parents and had segregation in F<sub>1</sub> or F<sub>2</sub> progenies

H.-j. Liu et al. (2011) studied shape and inner structure of stroma in *Cordyceps* using by cross section technique. The result showed that *C. sinensis* and *C. gunnii* found perithecia had ranged from elliptical to oval, embedded at the surface of the fertile portion of a stroma. In contract, *C. barnesii*, *C. liangshanesis* and *C. militaris* not found perithecia. The *C. barnesii* had outer cortex brown and inner cortex colourless. The *C. liangshanesis* had outer cortex brownish-yellow and inner cortex colourless. The *C. militaris* had colourless hyphae only.

T.-C. Wen et al. (2013) discovered new species of fungus called *C. sensu lato*. This species was identified by using by morphological technique, The *C. sensu lato* had cylindrical stroma, yellow with brown and the inner structure of mycelium sometimes abundantly and sometime negligent. The mycelium had gray and was found growing on Hepialidae host. This indicated that it can be described the difference among *Cordyceps* species.

## 2.10 Molecular markers used for genotyping fungi


A molecular marker is defined as a segment of DNA sequence, associated with a part of genome either linkage to gene of interest or target gene encoding a certain trait. The marker was widely used to evaluate DNA polymorphisms, patterns of heredity, genomic variations, evolutions, allele–allele linkages, and genetic map position in living organism (Joshi, Ranjekar, & Gupta, 1999). In *Cordyceps* species, most common markers used to determine genetic diversity were the random amplified polymorphic DNA (RAPD), developed to verify different species (Lam et al., 2015). The restriction fragment length polymorphism (RFLP) was also used to analyze the genetic relationship between two fungal species, such as *C. brongniartii* and *Beauveria brongniartii* (Sasaki, Miyamoto, Tamai, & Yajima, 2007). The inter-simple sequence repeat (ISSR) was evaluated the genetic diversity among eighteen *C. sisnensis* species in China (Liang et al., 2008). The internal transcribed spacer (ITS) sequence analysis was exploited to identify genetic difference between *C. sisnensis* from other related *Corrdyceps* species, such as *C. militaris* and *C. gunnii* (Chen et al., 2004). Additionally, the 28S rDNA were also used to study the genetic relationship and diversity among *Cordyceps* species (Artjariyasripong, Mitchell, HyweI-Jones, & Jones, 2001; G.-H. Sung, Spatafora, Zare, Hodge, & Gams, 2001). Of these markers, the ITS and 28S rDNA markers have been successfully used to identify nucleotide-sequence divergence in medicinal fungi, as previous publications in *Cordyceps* (Lam et al., 2015; Y. Li et al., 2013). For these, both markers were beneficially useful for describing genetic variation and structure of *C. militaris* populations. Therefore, the aim of this study was to classify genetic difference of *C. militaris* among species by using the ITS and 28S rDNA markers.

## 2.9 Definition of Internal transcribed spacer (ITS) and 28S ribosomal DNA

Internal transcribed spacer (ITS) refers to nucleotide sequences at the spacer DNA segments, located between the small-subunit ribosomal RNA (rRNA) and large-subunit rRNA genes in the chromosome or the corresponding to transcribed region in the polycistronic rDNA precursor (Peay, Kennedy, & Bruns, 2008). The ITS is especially useful for elucidating genetic relationships among species and closely related genus (Hao, Chen, Xiao, & Peng, 2010).

The 28S rDNA is the sequence of DNA that codes for ribosomal RNA, it was the first to be targeted for in detail studies of genetic variation in fungi (Simon, Binder, Adam, Hartig, & Ruis, 1992). The 28S have been useful to clarify the taxonomy among fungal phylum, Glomeromycota (Da Silva, Lumini, Maia, Bonfante, & Bianciotto, 2006). Sequence data from ribosomal sequences has represented an additional tool for the exact identification of fungal taxa, having low morphological divergence (Redecker, 2002). The 28S rDNA provides the genetic coding from which rDNA molecules are constructed.

The small subunits contain ITS1, 5.8S, ITS4, and 28S rDNA. The ITS1 is close to the 18S-rDNA border, and ITS 4 anneals to 28S rDNA near the ITS 4 border as shown in the Figure 5 (Lord, Kaplan, Shank, Kitts, & Elrod, 2002).



**Figure 5 Map positions of primer pairs (ITS1-ITS4 and LROR-LR7) used to amplify PCR products.**

Note: Primer ITS 1 is close to the 18S-rDNA border, and primer ITS 4 (5'-TCC TCC GCT TAT TGA TAT GC-3') anneals to 28S rDNA near the ITS 4 border.

## 2.10 Identification of *Cordyceps* using ITS and 28S rDNA markers

The previous researches, since 2000-2015, have been successfully used to the identification of *Cordyceps* species using ITS and 28S rDNA markers.

J. Kang, Liang, Liu, and Kong (2000) reported analysis of fungal DNA sequences, amplified by the 5.8S-rDNA and partial ITS2 markers. The result indicated that *C. gansuensis* and *C. sinensis* were conspecific (belonging to the same species).

Yue-Qin, Ning, Hui, and Liang-Hu (2002) studied genetic relationship between *C. sinensis* species, collected from different geographical regions of China.

The result found that the ITS sequences within *C. sinensis* have been high homologs, regardless of geographical origin. In contrast, *C. sinensis* was highly divergent from *C. militaris* and *C. gunnii*.

Z. Liu, Liang, Whalley, Liu, and Yao (2001) reported that DNA fragment of *C. sobolifera* was generated using a marker pair of ITS4 and ITS5. Resulting of the aligned DNA sequence between *C. sobolifera* and *Beauveria sobolifera*, collected from China, were not different.

Kinjo and Zang (2001) analyzed the phylogenetic relationships among the *C. sinensis* species from 11 geographic regions in Southwestern China, based on the sequences of ITS1-ITS2 and 28S rDNA regions. Both of markers indicated that the variations among different geographic regions were rather small.

Bae et al. (2002) reported that *C. militaris*, collected from Korea, Japan and China, were determined the genetic diversity by using ITS1 and ITS2 markers. All sequence analysis was low nucleotide divergence, ranged from 0.2% (1 bp) to 0.4% (2 bp). The genetic distance value was calculated by lower than 0.03. It was clearly shown that the nucleotide sequences at ITS region within *C. sinensis* were highly homologous, regardless of geographical origin.

Chen et al. (2004) report that three *Cordyceps* species (*C. sinensis* (synonym *O. sinensis*), *Hirsutella sinensis* and *Paecilomyces sinensis*) collected from Qingzang plateau, was analyze genetic variation by using the ITS1, ITS2, and 5.8S rDNA markers. The result show that the sequence comparison of *C. sinensis* was most closely related to *Hirsutella sinensis*, but clearly divergent from *Paecilomyces sinensis*.

(Kuo, Su, Yang, & Chen, 2005) reported that the ITS1, 5.8S rRNA, and ITS2 markers were used to construct a phylogenetic tree of 17 *Cordyceps* isolates. The result revealed that *C. sinensis* was less closely related to studied 15 *Cordyceps* species, but shared a closer relationship with *C. agriota*.

Wang, Zhang, Hu, Chen, and Qu (2008) investigated the genetic variation of *C. militaris* from different regions (Britain, China, Japan, Korea and Norway), and a PCR product was generated by using the ITS marker. The phylogenetic tree indicated that *C. militaris* was extremely small genetic variation ( $\leq 0.01$ , K2P model), which did not correlate to geographical origins. Moreover, this study indicated *C. militaris*



had the same distance level as *C. kyushuensis* and *C. roseostromata* collected from different regions.

Lam et al. (2015) reported that the ITS marker was developed to authenticate different genetic among *Cordyceps* species. The result found that twelve *C. sinensis* collected from geographical origin were highly homologous. In contrast, *C. sinensis* was high divergence from *C. gracilis*, *C. hawkesii* and *C. gunnii*.



## CHAPTER III

### MATERIALS AND METHODS

#### 3.1 Macroscopic studies

##### 1) Macroscopic characters of *C. militaris*

All *Cordyceps* sp. samples were collected from seven commercial farms in Thailand. The stromata of all samples was studied determination of morphological characteristics such as shape, size, colour and stromata.

##### 2) Mycelium growth

Individual fruiting bodies of *C. militaris* were collected from seven different commercial farms in Thailand. They were cut and plated on a potato dextrose agar (PDA) plate (containing 200 g/l potato, 20 g/l dextrose and 20 g/l agar), and cultured at 25°C in the dark for 7 days. The end of each mycelium was cut and plated on a new PDA plate. Colony diameter (cm) was measured at 3, 6, 9, 12, 15, 18 and 21 days of culture. Mean comparison between *C. militaris* strains were calculated by were compared Duncan's Multiple Range Test (DMRT).

#### 3.2 Microscopic studies

##### 1) Cross section of stromata

The stroma was cut using cross section techniques and stained with lactophenol cotton blue. Morphological characteristics such as surface color, mycelium color and perithecium formation were observed under a microscope (Olympus CH30RF200, Japan) and photographed.



## **2) Spore shooting**

The spores from the stromata were allowed to shoot upwards onto a cover-slip, placed on the middle PDA medium plate, and incubated at room temperature for 24 hours in dark condition. The cover-slip was removed, and then placed upside down on a new slide, containing a drop of lactophenol cotton blue. Morphology of spores was assessed under the microscope.

## **3) Slide culture technique**

Slide culture technique is a valuable technique to study fungal morphology in hyphae and conidia state. Firstly, the mycelium of *C. militaris* was cultivated on PDA at 25°C under dark condition for 7 days. Meanwhile, the slide culture technique was set up as following steps the V shaped glass rod was placed in sterilized petri dishes, the slide was put on the V shaped glass rod, a small agar piece (0.5x0.5x0.5 mm<sup>3</sup>) was placed on the coverslip, the mycelium (cultivated on PDA) was picked up using a needle to inoculate on the four side of the face-centered square of agar. The coverslip was carefully placed over the agar block; sterile distilled water (5 ml) was poured into the petri dishes. Finally, the slide culture set was incubated at 25°C under dark conditions for 14 days. At the end of experiment, the agar piece was removed, and the mycelia were grown on the slide and coverslip, and then produced conidia. It was stained by lactophenol cotton blue and observed using a compound microscope.

## **4) SEM Study**

Mycelium was individually cultured on PDA medium at 25°C under dark condition for 14 days. The growing mycelium was cut with sterile scalpels into small pieces (1x1 cm) and fixed in 2.5% (v/v) glutaraldehyde. It was washed in 0.1 M phosphate buffer for 15 minutes repeated twice, and washed with 1% osmium tetroxide for 1 hour. The mycelium was then dehydrated through a series of 30, 50, 70, 90 and 100% ethyl alcohol for 30 minutes, respectively. Finally, the mycelium was dehydrated in acetone for 30 minutes before mounting on an aluminum stub and coating with gold-palladium alloy. Spore structures were examined under a scanning electron microscope (JEOL LV 5910, Japan).

### 3.3 Genomic DNA (gDNA) extraction and PCR amplification

The medium, grown on PDA for 3-7 days, was collected for genomic DNA (gDNA) extraction using InstaGene Matrix (Bio-Rad, United States) according to the manufacturer's instructions.

Two primer pairs as ITS primer pair (ITS1 and ITS4) and 28S rDNA primer pair (LR0R and LR7) were used in this study; their sequences are shown in Table 1. The primers of ITS1 and ITS4 were located on the 3' end of 18S-rDNA border and the end of 28S rDNA border, respectively (Figure. 5). The ITS primer pair was used to amplify the DNA fragment to flank the entire ITS region and intervening 5.8S subunit (Figure. 5). The PCR reaction was carried out with 20 ng of gDNA as the template in a 30 µl reaction mixture using an EF-*Taq* (SolGent, Korea). The PCR condition was performed with the initial denaturation as follows: activation of *Taq* polymerase at 95°C for 2 minutes, followed by 35 cycles of 95°C for 1 minute, 55°C and 72°C for 1 minute each, and a final extension at 72°C for 10 minutes.


Meanwhile, the LR0R and LR7 primers were used for the DNA fragment at the partial 28S rDNA region. The amplified products were purified using a multiscreen filter plate (Millipore Corp., Bedford, MA, United States). Purified DNA samples were performed using a PRISM BigDye Terminator v3.1 Cycle Sequencing Kit (Applied Biosystems, Foster City, Canada) containing the extension products added to Hi-Di formamide. The mixture was incubated at 95°C for 5 minutes, followed by 5 minutes on ice, and then sequenced and analyzed using an ABI Prism 3730XL DNA analyzer (Applied Biosystems, Foster City, Canada).

### 3.4 DNA sequence alignment

Partial nucleotide-sequence reads were approximately 548 and 1223 nucleotides, amplified by ITS and 28S rDNA primer pairs, respectively. Both sequence reads were assembled using Bioedit Analysis software (Hall, 1999) and compared to known sequences of ITS and 28S rDNA regions of fungi published from the NCBI database using CLUSTAL W (European Bioinformatics Institute (EMBL-EBI), United Kingdom).

**Table 2 Primers used to amplify fungal 28S rDNA genes and ITS region sequences**

<b>Primer</b>	<b>Sequence (5'→3')</b>	<b>Reference</b>
<b>28S rDNA region</b>		
LROR	ACCCGCTGAACTTAAGC	(Bunyard, Nicholson, & Royse, 1994)
LR7	TACTACCACCAAGATCT	(Bunyard et al., 1994)
<b>ITS region</b>		
ITS1	TCCGTAGGTGAACCTGCGG	(White, Bruns, Lee, & Taylor, 1990)
ITS4	TCCTCCGCTTATTGATATGC	(White et al., 1990)



## CHAPTER IV

### RESULTS AND DISCUSSION

#### 4.1 Macroscopic studies

##### 1) Macroscopic characters of *C. militaris*

In this study, specimens of *C. militaris* have been collected from seven commercial farms in Thailand. Morphological characteristics (shape, size and colour of the stromata, including growth rate and colour of mycelium) of *C. militaris* (*Cmi1*-*Cmi7*). Seven isolates were examined through the macroscopic (naked eyes) and microscopic techniques. All studied isolates had only a single golden yellow stroma (Figure 6). The stroma of *Cmi4* had the shortest and widest stipe, compared to other isolates, width 1.5-2.5 cm length 2-4 cm. The *Cmi7* had the longest stroma stipe (width 0.3-0.4 cm length 5.2-5.5 cm), and its head was indented. Moreover, the stromata of *Cmi1*, *Cmi2* and *Cmi3* had cylindrical stipes (width 0.3-0.8 cm and length 4-6 cm), and head of stoma were spherical (Figure 6). Resulting of different stromatal characters among studied *C. militaris* would be caused by various culture condition effects, which were supported by many published reports that morphological changes of *C. militaris* relied on pH (J.-P. Park, Kim, Hwang, & Yun, 2001) and (G.-H. Sung et al., 2011), temperature (J.-M. Sung, Choi, Shrestha, & Park, 2002), humidity (Tapingkae, 2012), light intensity (Sato & Shimazu, 2002) and (G.-H. Sung et al., 2011), carbon, nitrogen and mineral sources culture media (G.-H. Sung et al., 2011), strains and culture methods (Lee et al., 2013). Moreover, different cultivation times may affect morphological changes, which had been reported by G. Zhang and Liang (2013) and J.-M. Sung et al. (2002) who respectively found that the suitable harvesting times for *C. militaris* 45 days and 50 days.




**Figure 6 Morphological characteristic of stromata among *C. militaris* isolates (Cmi1- Cmi7)**

## **2) Mycelium growth**

Pathania et al. (2015) reported that mycelium growth of *C. militaris* cultured on PDA (Potato dextrose agar) medium was not significantly different from a PDA medium. In this experiment, the PDA medium was used to study mycelium growth of seven isolates (*C. militaris*) because it was a simple and inexpensive preparation. All isolates of *C. militaris* were cultured on the PDA medium in a modified incubator at 25°C in the dark for 21 days, and mycelium-growth rates were examined every three days. The result revealed that the mycelial growth rates were divided into two groups. Firstly, highest growth rate (*Cmi1*, *Cmi2*, *Cmi3*, *Cmi5*, *Cmi6*, and *Cmi7*) had a colony diameter ranged from  $1.90 \pm 0.17$  to  $8.50 \pm 0.00$  cm. Secondly, *Cmi4* had the lowest growth rate (Figure 7) and gave colony diameter ranged from  $1.78 \pm 0.09$  to  $6.94 \pm 0.14$  cm) (Table 4). Meanwhile, seven isolates of *C. militaris* were cultured on the



PDA medium in modified incubator at 25°C under dark conditions for 21 days, the growth characteristics were determined every 7 days until 21 days. The results found that the mycelial characteristics of six isolates (*Cmi1*, *Cmi2*, *Cmi3*, *Cmi5*, *Cmi6*, and *Cmi7*) were abundant mycelial density, rough and semi-cottony mycelial texture, and yellowish colony (Table 3 and Table 5). Only the mycelial characteristic of *Cmi1* was formed stroma on the PDA medium (Figure 7). Moreover, *Cmi4* mycelium was compact formed densely layers, and dark yellow in middle and white at the edges on PDA (Figure 7). These results indicate that mycelial growth rates and the mycelial characteristics of individually studied isolates were significantly varied, which may depend on different cultivation times (Lee et al., 2013; Shrestha et al., 2006), and different strains (Lee et al., 2013).



**Figure 7** Mycelial growth rates and characteristics among seven *C. militaris* isolates (*Cmi1* to *Cmi7*) cultured on PDA medium for 3, 6, 9, 12, 15, 18 and 21 days

**Table 3 Mycelial growth characteristics of seven *C. militaris* isolates (*Cmi1-Cmi7*) on PDA**

isolates	Mycelial colony					
	7 days		14 days		21 days	
	Mycelial density	Mycelial texture	Mycelial density	Mycelial texture	Mycelial density	Mycelial texture
<i>Cmi1</i>	+ to ++	R to SC	++	F to C	++	F to C
<i>Cmi2</i>	+, ++	R to SC	++	R to SC	++	R to SC
<i>Cmi3</i>	+, ++	R to SC	++	R to SC	++	R to SC
<i>Cmi4</i>	+, ++	R to SC	++	R to SC	++	R to SC
<i>Cmi5</i>	+, ++	F to SC	++	F to C	++	F to C
<i>Cmi6</i>	+, ++	R to SC	++	R to SC	++	R to SC
<i>Cmi7</i>	+, ++	R to SC	++	R to C	++	R to C

Note: + indicates moderate density; ++ indicates abundant density, F indicates flat mycelial texture, R indicates rough mycelial texture; SC indicates semi-cottony mycelial texture; and C indicates cottony mycelial texture

**Table 4 Colony diameter of seven *C. militaris* (*Cmi1- Cmi7*) isolates on PDA medium**

strain	Colony diameter measured of <i>C. militaris</i> (cm)						
	3 days	6 days	9 days	12 days	15 days	18 days	21 days
<i>Cmi1</i>	1.39±0.11 <sup>c</sup>	2.46±0.05 <sup>d</sup>	4.14±0.07 <sup>b</sup>	5.93±0.52 <sup>bc</sup>	5.86±0.40 <sup>bc</sup>	7.96±0.03 <sup>ab</sup>	8.50±0.00 <sup>a</sup>
<i>Cmi2</i>	1.90±0.05 <sup>a</sup>	3.25±0.15 <sup>a</sup>	4.30±0.18 <sup>b</sup>	5.48±0.24 <sup>c</sup>	7.13±0.85 <sup>ab</sup>	7.05±0.40 <sup>c</sup>	8.50±0.00 <sup>a</sup>
<i>Cmi3</i>	1.58±0.08 <sup>c</sup>	3.31±0.07 <sup>a</sup>	5.08±0.10 <sup>a</sup>	6.45±0.22 <sup>ab</sup>	7.23±0.49 <sup>ab</sup>	8.50±0.00 <sup>a</sup>	8.50±0.00 <sup>a</sup>
<i>Cmi4</i>	1.78±0.09 <sup>ab</sup>	2.73±0.06 <sup>cd</sup>	3.61±0.07 <sup>c</sup>	4.59±0.24 <sup>d</sup>	5.50±0.28 <sup>c</sup>	6.33±0.12 <sup>d</sup>	6.94±0.14 <sup>b</sup>
<i>Cmi5</i>	1.86±0.01 <sup>ab</sup>	3.08±0.36 <sup>bc</sup>	4.36±0.19 <sup>b</sup>	5.70±0.20 <sup>bc</sup>	6.88±0.15 <sup>ab</sup>	7.81±0.20 <sup>b</sup>	8.50±0.00 <sup>a</sup>
<i>Cmi6</i>	1.58±0.08 <sup>bc</sup>	2.78±0.37 <sup>cd</sup>	4.33±0.07 <sup>b</sup>	6.26±0.10 <sup>bc</sup>	7.36±0.06 <sup>a</sup>	8.45±0.05 <sup>a</sup>	8.50±0.00 <sup>a</sup>
<i>Cmi7</i>	1.97±0.12 <sup>a</sup>	2.46±0.16 <sup>ab</sup>	4.86±0.88 <sup>a</sup>	6.85±0.07 <sup>a</sup>	7.60±0.15 <sup>a</sup>	8.20±0.15 <sup>b</sup>	8.50±0.00 <sup>a</sup>

Note: colony diameter value (mm) represents by mean ± se, was measured from all *C. militaris* isolates (*Cmi1 -Cmi7*) with triplicates. The same letters in each column indicates non-significant difference of mean values, calculated by using DMRT method.



**Table 5 Colony pigmentation of *C. militaris* isolates (*Cmi1- Cmi7*) on PDA medium**


Strain	Colony pigmentation during					
	6 days		12 days		18 days	
	Margin	Center	Margin	Center	Margin	Center
<i>Cmi1</i>	Y	YW	W	YW to Y	W	Y
<i>Cmi2</i>	W to PY	YW	W to YW	Y	W to YW	Y
<i>Cmi3</i>	W to PY	PY	W to YW	W to PY	W to YW	W to YW
<i>Cmi4</i>	W to YW	MY	W to MY	MY	W	W to MY
<i>Cmi5</i>	W	W	W to YW	YW	W	W to Y
<i>Cmi6</i>	W	W	W to YW	YW	W to YW	W to YW
<i>Cmi7</i>	W to PY	W	W to Y	W	W to YW	W to YW

Note: W indicates white; Y indicates yellow; YW indicates yellowish white; PY indicates pale yellow; and MY indicates mustard yellow.

## 4.2 Microscopic characteristics studies

### 1) Microscopic characteristics of stromata among *C. militaris*


The outer surface of the seven *C. militaris* isolates (Figure 8) found that most studied isolates had long, smooth, golden yellow stipe and round-surface stromata, except for the *Cmi4* had a short stipe and sharp cap. These were agreed with Zhang et al., 2013 reported that the cultivated stromata of *C. militaris* was clavate with smooth shaft and orange stipe. Moreover, the *Cmi5* and *Cmi6* had a rugged surface. These may be possible that they were different strains, ages of stroma and condition of culture media (Lee et al., 2013).



**Figure 8 The outer surface characteristics of stromata among seven *C. militaris* isolates observed under Stereo microscope**

## **2) Cross section of stromata among *C. militaris***

Cross sections of individual stroma morphologies of the seven *C. militaris* isolates were examined by staining with lactophenol cotton blue. All isolates showed similar stroma characteristics with smooth yellow gold color outer cortex surfaces, and inner cortexes containing colorless hyphae without a perithecia (Figure 9). These results indicated that the presence or absence of perithecia might be beneficially used to evaluate the morphological diversity of *C. militaris*. As supported by previous publications, some *C. militaris* species showed the presence of perithecia (Xiong, Xia, Zheng, Shi, & Wang, 2010) but other species did not have perithecia (H.-j. Liu et al., 2011) in the cortex region of the stroma. In contrast, all *C. militaris* species had a smooth outer cortex surface and colorless hyphae at the inner cortex (H.-j. Liu et al., 2011).




**Figure 9 Evaluation of stromal characteristics among seven *C. militaris* isolates (*Cmi1* to *Cmi7*) by cross section technique under a compound microscope.**

Note: random stroma (A) were cross sectioned to examine microscopic characteristics (B); the scale bar indicates 10X (=60  $\mu$ m)

### 3) Spore shooting technique

Spore shooting technique is basically used to study spore characteristics among fungal species, based on a single spore (Strobel et al., 1996). In this research, study of spores from stromata *C. militaris* isolates were discharged on cover slip and their morphology was determined under a microscope. The result revealed that all isolates had cylindrical, globose and formed oval spores (Figure 10). These spores were similar to result from the study of Shrestha et al. (2005) who reported that freshly discharged ascospore of *C. militaris* in sterile water showed spores oval and oblong in shape.

Moreover, it was found that the discharged spore produced germ tube after 24 hours, in agreement with Tulasne (1865) found that the spore of *C. militaris* usually germinate in a short time period.




**Figure 10** Spore production of *C. militaris* Cmi1-Cmi7 isolates by spore shooting

Note: bar indicates 40X (=20 µm)

#### 4) Slide culture technique

To examine the hyphae germination and conidial formation of seven different *C. militaris* isolates (Cmi1- Cmi7), an individual mycelium was cultivated on PDA medium at 25°C under dark conditions for 14 days by using slide culture technique. The result found that all isolates had similar hyphae morphology and conidial formation. Most isolates produced long hyphae, formed several branches and first-formed conidia as ovoid, pyriform or cylindrical and slim heads. However, some conidia formed short chains, and had a globose head (Figure 11). It has been reported that specimens of different *C. militaris* strains formed branched germinating hyphae, produced conidia in short chains, globose head and slimy head, first-formed conidia were ovoid, pyriform or cylindrical, with the succeeding conidia being globose, nearly subglobose or ellipsoid (Shrestha et al., 2005).





**Figure 11 Spore germination of different isolates of *C. militaris* Cmi1-Cmi7 by slide culture technique for 14 days, (A) Conidia formed in short chains (B) Conidia formed in globose head**


Note: bar indicates A = 100X, B = 100X (=0.06 μm)

### **5) Observation of conidial formation by Scanning Electron Microscopy (SEM)**

Morphologies of the conidia of seven *C. militaris* isolates (Cmi1 to Cmi7) were determined using SEM. Results showed that most conidia were globose and oval shaped which germinated to produce long hyphae. Most conidia were formed at the end of the hyphae as single globe heads but rare groups were globose in short chains. A few conidia formed as oval shapes (Figure 12).

As previously reported, hyphae of *C. militaris* (EFCC 11255) produce various forms of conidia, either as a single globose head or groups of globose short chains

(Shrestha et al., 2005). Moreover, conidia of *C. militaris* were presented as various shapes including globose (Tulasne, 1865) cylindrical and spherical (Brown & Smith, 1957; De Bary, 1867), almost spherical (Pettit, 1895), and oval (Petch, 1936).



**Figure 12** Scanning electron micrographs of conidial hyphae of *C. militaris* Cmi1-Cmi7 14 days after culture on PDA (Scales: A = 1  $\mu$ m, B = 5  $\mu$ m)

#### 4.3 DNA sequencing

Both the internal transcribed spacer (ITS) and 28S rDNA regions are highly divergent in nucleotide sequences and play a vital role in evolution among fungal species (Lam et al., 2015). Therefore, these areas were used to assess genetic variation among the seven isolates of *C. militaris* based on PCR assay. Two ITS primer pairs (ITS1 and ITS4) were used to amplify the DNA fragment flanking the entire ITS region and intervening 5.8S subunit, giving an ITS-PCR-fragment (548 bp long). Nucleotide sequences from all the studied isolates gave 100% similarity, indicating



that the seven isolates of *C. militaris* did not have genetic distance at the ITS. This distance values of the ITS region among the seven isolates were slightly different when compared to *C. militaris* isolates NBRC100741 (Figure 13) (Schoch et al., 2012). Only one nucleotide (adenine, A) at position 93 of the ITS region of each studied strain was substituted by guanine (G) of NBRC100741 strain (Schoch et al., 2012) (Figure 13). The nucleotide substitution might be caused by evolutionary effect (Y.-J. Park & Min, 2005), and provide information for phylogenetic reconstruction (Watanabe et al., 2011) fungal species.

Meanwhile, two 28S rDNA primer pairs (LROR and LR7) were used to amplify the DNA fragment at the 28S rDNA region, giving a 28S rDNA-PCR-fragment (1223 bp long). Nucleotide sequence reads of both the ITS-PCR-fragment and the 28S rDNA-PCR-fragment from individual isolates were aligned using the ClustalW program and then compared to their known fungal sequences published from the NCBI database. Also, nucleotide sequences from all the studied isolates gave 100% 28S rDNA (Figure 16). This distance values of the 28S rDNA regions among the seven isolates were slightly different when compared to *C. militaris* strains JM0807 (Zhong, Peng, Qi, Lei, & Liu, 2010). Furthermore, a nucleotide at position 136 in the 28S rDNA region of *C. militaris* JM0807 strain was inserted by a thymine base (T) compared to all studied strains, as previously reported by Zhong et al. (2010) (Figure 14). The nucleotide insertions can be correlated with the evolutionary distance among fungal species (Y.-J. Park & Min, 2005).

However, nucleotide sequences, located on ITS and 28S region of all studied *C. militaris* isolates were compared to other *C. militaris* strains by using Blast data program. The results found that these comparisons had 99-100% and 99% the percentages of identity at ITS (Table 6) and 28S (Table 7), respectively. These confirmed that all studied isolates were truly *C. militaris* species.

In summary, comparative analysis of nucleotide sequences both ITS and 28SrDNA were clearly indicated that there were not genetic variation among all studied *C. militaris* isolates. However, they exhibited only one nucleotide substitution in ITS and one nucleotide insertion in 28SrDNA, compared to *C. militaris* strains, NBRC100741 or JM0807, respectively.

<b>Samples</b>	1	AAGTTCAGCGGGTATTCCTACCTGATCCGAGGTCAACGTTTCAGAGTTGGGCGTTTACGG	60
<b>Ref</b>	785	AAGTTCAGCGGGTATTCCTACCTGATCCGAGGTCAACGTTTCAGAGTTGGGCGTTTACGG	726
<b>Samples</b>	61	CGTGGCCACGTCGGGTTCCCGGTGCGAGTTGGAGTACTACGCAGAGGTCGCCGCGGACGG	120
<b>Ref</b>	725	CGTGGCCACGTCGGGTTCCCGGTGCGAGTTGGAGTACTACGCAGAGGTCGCCGCGGACGG	666
<b>Samples</b>	121	GCCGCCACTTCATTTCCGGGGCGGCGGTGTGCTGCCGGTCCCCAACGCCGACATCCCCCA	180
<b>Ref</b>	665	GCCGCCACTTCATTTCCGGGGCGGCGGTGTGCTGCCGGTCCCCAACGCCGACATCCCCCA	606
<b>Samples</b>	181	GGGGACGTCGAGGGTTGAAATGACGCTCGAACAGGCATGCCCGCCAGAATGCTGGCGGGC	240
<b>Ref</b>	605	GGGGACGTCGAGGGTTGAAATGACGCTCGAACAGGCATGCCCGCCAGAATGCTGGCGGGC	546
<b>Samples</b>	241	GCAATGTGCGTTCAAAGATTTCGATGATTCACTGAATTCTGCAATTCACATTACTTATCGC	300
<b>Ref</b>	545	GCAATGTGCGTTCAAAGATTTCGATGATTCACTGAATTCTGCAATTCACATTACTTATCGC	486
<b>Samples</b>	301	ATTTTCGCTGCGTTCTTCATCGATGCCAGAGCCAAGAGATCCGTTGTTGAAAGTTTGATT	360
<b>Ref</b>	485	ATTTTCGCTGCGTTCTTCATCGATGCCAGAGCCAAGAGATCCGTTGTTGAAAGTTTGATT	426
<b>Samples</b>	361	CATTTGTTTTGCCTTGCGGCGGATTTCAGAAAACTGGTAGATACAGTGTGTTGGGGCCCC	420
<b>Ref</b>	425	CATTTGTTTTGCCTTGCGGCGGATTTCAGAAAACTGGTAGATACAGTGTGTTGGGGCCCC	366
<b>Samples</b>	421	GACGGCCGCCGCCAGGCCCGCTCCAGGCGCTGGGCGAGTCCGCCGAAGCAACGATAGG	480
<b>Ref</b>	365	GACGGCCGCCGCCAGGCCCGCTCCAGGCGCTGGGCGAGTCCGCCGAAGCAACGATAGG	306
<b>Samples</b>	481	TATGTTCAAAAGGGTTGGGAGTTGGAAAACTCGTTAATGATCCCTCCGCTGGTTCACCA	540
<b>Ref</b>	305	TATGTTCAAAAGGGTTGGGAGTTGGAAAACTCGTTAATGATCCCTCCGCTGGTTCACCA	246
<b>Samples</b>	541	ACGGAGAC	548
<b>Ref</b>	245	ACGGAGAC	238

**Figure 13 Comparative analysis of nucleotide sequences between seven studied *C. militar* isolates and *C. militar* NBRC100741 strain using ClastaW program**

Note: samples indicate seven studied *C. militar* isolates. Ref indicates *C. militar* NBRC100741 strain, yellow lighted letter indicates the nucleotide substitution in ITS region, and number indicates position of nucleotide sequences.

**Table 6 Blast results of the ITS nucleotide sequences generated from studied *C. militaris* isolates using nucleotide blast (NCBI)**

Description	Query cover	Identities	Accession
<i>C. militaris</i> isolate WU-CMH	100%	100%	KM197165.1
<i>C. militaris</i> strain NBRC 30377	99%	100%	JN943300.1
<i>C. militaris</i> strain NBRC 100741	100%	99%	JN943437.1
<i>C. militaris</i> strain NBRC 9787	100%	99%	JN943433.1
<i>C. militaris</i> strain BCRC33736	100%	99%	HQ591387.1
<i>C. militaris</i> strain SPNU1002	96%	99%	KY407774.1
<i>C. militaris</i> strain KACC50001	96%	99%	KY407771.1
<i>C. militaris</i> strain KACC44454	96%	99%	KY407766.1
<i>C. militaris</i> strain KACC40226	96%	99%	KY407760.1

Note: Query cover is the percentage of sequences from seven *C. militaris* isolates, aligned to a sequence in each accession of *C. militaris* strain.

Identity is the percentage of sequences from seven *C. militaris* isolates which is highly significant match to reference sequences.

<b>Samples</b>	1	GAAATCTGGCCCCCGGTCCGAGTTGTAATTTGTAGAGGATGCTTTTGGCGAGGTGCCTT	60
<b>Ref</b>	95	GAAATCTGGCCCCCGGTCCGAGTTGTAATTTGTAGAGGATGCTTTTGGCGAGGTGCCTT	154
<b>Samples</b>	61	CCGAGTTCCTTGAACGGGACGCCATAGAGGGTGAGAGCCCCGTCTGGTCGGACACCGAG	120
<b>Ref</b>	155	CCGAGTTCCTTGAACGGGACGCCATAGAGGGTGAGAGCCCCGTCTGGTCGGACACCGAG	214
<b>Samples</b>	121	CCTCTGTAAAGCTCC-TTCGACGAGTCGAGTAGTTTGGGAATGCTGCTCAAAATGGGAGG	179
<b>Ref</b>	215	CCTCTGTAAAGCTCC-TTCGACGAGTCGAGTAGTTTGGGAATGCTGCTCAAAATGGGAGG	274
<b>Samples</b>	180	TATATGTCTTCTAAAGCTAAATATTGGCCAGAGACCGATAGCGCACAAAGTAGAGTGATCG	239
<b>Ref</b>	275	TATATGTCTTCTAAAGCTAAATATTGGCCAGAGACCGATAGCGCACAAAGTAGAGTGATCG	334
<b>Samples</b>	240	AAAGATGAAAAGCACTTTGAAAAGAGGGTTAAAAAGTACGTGAAATTGTTGAAAGGGAAG	299
<b>Ref</b>	335	AAAGATGAAAAGCACTTTGAAAAGAGGGTTAAAAAGTACGTGAAATTGTTGAAAGGGAAG	394
<b>Samples</b>	300	CGCCTATGACCAGACTTGGGCCCGGTGAATCACCAGCGTTCTCGCTGGTGCCTTCGCC	359
<b>Ref</b>	395	CGCCTATGACCAGACTTGGGCCCGGTGAATCACCAGCGTTCTCGCTGGTGCCTTCGCC	454
<b>Samples</b>	360	GGGCACAGGCCAGCATCAGTTTGGCGCGGGGAGAAAGGCTTCGGGAATGTGGCTCCCT	419
<b>Ref</b>	455	GGGCACAGGCCAGCATCAGTTTGGCGCGGGGAGAAAGGCTTCGGGAATGTGGCTCCCT	514
<b>Samples</b>	420	GGGAGTGTTATAGCCCGTTGCGCAATACCCTGCGCTGGACTGAGGTACGCGCATTCGCAA	479
<b>Ref</b>	515	GGGAGTGTTATAGCCCGTTGCGCAATACCCTGCGCTGGACTGAGGTACGCGCATTCGCAA	574
<b>Samples</b>	480	GGATGCTGGCGTAATGGTCATCAGCGACCCGTCTTGAAACACGGACCAAGGAGTCGTCTT	539
<b>Ref</b>	575	GGATGCTGGCGTAATGGTCATCAGCGACCCGTCTTGAAACACGGACCAAGGAGTCGTCTT	634
<b>Samples</b>	540	CGTATGCGAGTGTTCCGGGTGTCAAACCCCTACGCGTAATGAAAGTGAACGCAGGTGAGAG	599
<b>Ref</b>	635	CGTATGCGAGTGTTCCGGGTGTCAAACCCCTACGCGTAATGAAAGTGAACGCAGGTGAGAG	694
<b>Samples</b>	600	CTTCGGCGCATCATCGACCGATCCTGATGTTCTCGGATGGATTGAGTAAGAGCATACGG	659
<b>Ref</b>	695	CTTCGGCGCATCATCGACCGATCCTGATGTTCTCGGATGGATTGAGTAAGAGCATACGG	754
<b>Samples</b>	660	GGCCGGACCCGAAAGAAGGTGAACTATGCCTGTATAGGGTGAAGCCAGAGGAAACTCTGG	719
<b>Ref</b>	755	GGCCGGACCCGAAAGAAGGTGAACTATGCCTGTATAGGGTGAAGCCAGAGGAAACTCTGG	814
<b>Samples</b>	720	TGGAGGCTCGCAGCGGTTCTGACGTGCGAATCGATCGTCAAATATGGGCATGGGGGCGAA	779
<b>Ref</b>	815	TGGAGGCTCGCAGCGGTTCTGACGTGCGAATCGATCGTCAAATATGGGCATGGGGGCGAA	874
<b>Samples</b>	780	AGACTAATCGAACCTTCTAGTAGCTGGTTTCCGCCGAAGTTTCCCTCAGGATAGCAGTGT	839
<b>Ref</b>	875	AGACTAATCGAACCTTCTAGTAGCTGGTTTCCGCCGAAGTTTCCCTCAGGATAGCAGTGT	934
<b>Samples</b>	840	TGAACTCAGTTTATGAGGTAAAGCGAATGATTAGGGACTCGGGGCGCTATATTGCCTT	899
<b>Ref</b>	935	TGAACTCAGTTTATGAGGTAAAGCGAATGATTAGGGACTCGGGGCGCTATATTGCCTT	994

<b>Samples</b>	900	CATCCATTCTCAAACCTTTAAATATGTAAGAAGCCCTTGTTACTTGATTGAACGTGGGCAT	959
<b>Ref</b>	995	CATCCATTCTCAAACCTTTAAATATGTAAGAAGCCCTTGTTACTTGATTGAACGTGGGCAT	1054
<b>Samples</b>	960	TCGAATGTATCAACACTAGTGGGCCATTTTTGGTAAGCAGAACTGGCGATGCGGGATGAA	1019
<b>Ref</b>	1055	TCGAATGTATCAACACTAGTGGGCCATTTTTGGTAAGCAGAACTGGCGATGCGGGATGAA	1114
<b>Samples</b>	1020	CCGAACGCGAGGTTAAGGTGCCGGAGTGGACGCTCATCAGACACCACAAAAGGTGTTAGT	1079
<b>Ref</b>	1115	CCGAACGCGAGGTTAAGGTGCCGGAGTGGACGCTCATCAGACACCACAAAAGGTGTTAGT	1174
<b>Samples</b>	1080	ACATCTTGACAGCAGGACGGTGGCCATGGAAGTCGGAATCCGCTAAGGACTGTGTAACAA	1139
<b>Ref</b>	1175	ACATCTTGACAGCAGGACGGTGGCCATGGAAGTCGGAATCCGCTAAGGACTGTGTAACAA	1234
<b>Samples</b>	1140	CTCACCTGCCGAATGTACTAGCCCTGAAAATGGATGGCGCTCAAGCGTCCCACCCATACC	1199
<b>Ref</b>	1235	CTCACCTGCCGAATGTACTAGCCCTGAAAATGGATGGCGCTCAAGCGTCCCACCCATACC	1294
<b>Samples</b>	1200	TCGCCCTCAGGGTAGAAACGAAGC	1223
<b>Ref</b>	1295	TCGCCCTCAGGGTAGAAACGAAGC	1318

**Figure 14 Comparative analysis of nucleotide sequences between seven studied *C. militaris* isolates and *C. militaris* JM0807 strain using ClastaW program**

Note: samples indicate seven studied *C. militaris* strains, Ref indicates *C. militaris* JM0807 strain, yellow lighted letter indicates the nucleotide insertion in 28S rDNA region, and number indicates position of nucleotide sequences

**Table 7** Blast results of the 28S rDNA nucleotide sequences generated from studied *C. militaris* isolates using nucleotide blast (NCBI)

Description	Query cover	Identities	Accession
<i>C. militaris</i> strain ATCC 34164	100%	99%	CP023322.1
<i>C. militaris</i> strain JM0807	100%	99%	JN411084.1

Note: Query cover is the percentage of sequences from seven *C. militaris* isolates, aligned to a sequence in each accession of *C. militaris* strain.

Identity is the percentage of sequences from seven *C. militaris* isolates which is highly significant match to reference sequences.

## CHAPTER V

### CONCLUSION

Seven commercial isolates of *C. militaris* (*Cmi1*, *Cmi2*, *Cmi3*, *Cmi4*, *Cmi5*, *Cmi6* and *Cmi7*) collected in Thailand were investigated for morphological study using various macroscopic and microscopic methods.

In macroscopic studies, the results concluded that most isolates had cylindrical shapes and golden yellow of stromata. Moreover, they had the highest mycelial growth rate and abundant mycelial density with rough and semi-cottony mycelial texture, and yellowish colony. Of these, *Cmi4* had the lowest growth rate, and its mycelium was compact, formed of dense layers, and dark yellow on PDA.

In microscopic studies using a cross section technique, the stromal characteristics revealed that all isolates were golden yellow at the edge, and colourless hyphae. The spore characteristics, investigated by using the spore shooting technique, revealed that most conidiospores were globular shape. Conidial formation was studied using slide culture and SEM techniques. Results determined that most spores were globose and oval shaped; moreover, these germinated and produced long hyphae. At the ends of the hyphae, most conidia were formed as single globe heads but rarely as global groups in short chains. A few conidia were formed with an oval shape.


Meanwhile, all studied *C. militaris* isolates were evaluated genetic variation by using nucleotide sequence analysis of ITS and 28S rDNA regions. Results concluded that all studied isolates showed no genetic variation. However, they had one nucleotide substitution in the ITS region and one nucleotide insertion in the 28S rDNA region, compared to *C. militaris* NBRC100741 or JM0807 strains, respectively.



## Suggestion

These results were clearly elucidated that the macroscopic characteristics (such as stroma characteristic, mycelial growth rate, and mycelial pigmentation) were basically significant tools for fungal classification, including *C. militaris* isolates of this studied.

However, the microscopic characteristics and molecular markers (in case ITS and 28S rDNA) have been still not found differences. Therefore, further research is necessary to collect more samples (*C. militaris*) from different commercial sources in Thailand to examine morphological techniques and other molecular markers such as random amplification of polymorphic (RAPD), terminal restriction fragment (TRF) and amplified ribosomal DNA restriction analysis (ARDRA). Furthermore, it should be study quantity and quality of bioactive compound (cordycepin, adenosine) in *C. militaris*.



## REFERENCES



- Albini, M. (2017). *Fungal Biogenic Patina: Optimization of an Innovative Conservation Treatment for Copper-based Artefacts*: Université de Neuchâtel, Faculté des sciences.
- Alexopoulos, C., Mims, C., & Blackwell, M. (1996). *Introductory mycology*. 1996. New York.
- Artjariyasripong, S., Mitchell, J. I., Hywel-Jones, N. L., & Jones, E. B. G. (2001). Relationship of the genus *Cordyceps* and related genera, based on parsimony and spectral analysis of partial 18S and 28S ribosomal gene sequences. *Mycoscience*, 42(6), 503-517.
- Avtonomova, A., Krasnopolskaya, L., Shuktueva, M., Isakova, E., & Bukhman, V. (2015). Assessment of antitumor effect of submerged culture of *Ophiocordyceps sinensis* and *Cordyceps militaris*. *Antibiotiki i khimioterapiia= Antibiotics and chemotherapy [sic]*, 60(7-8), 14-17.
- Bae, J.-S., Jin, B.-R., Lee, H.-O., Park, E.-J., Yu, L., & Lee, S.-M. (2002). Determination of the Ribosomal DNA Internal Transcribed Spacers and 5.8S rDNA Sequences of *Cordyceps* Species. *International Journal of Industrial Entomology*, 5(1), 85-91.
- Bhandari, A. K., Negi, J., Bisht, V., Rana, C., Bharti, M., & Singh, N. (2010). Chemical constituent, inorganic elements and properties of *Cordyceps sinensis*—a review. *Nature and Science*, 8(9), 253-256.
- Brown, A. H., & Smith, G. (1957). The genus *Paecilomyces* Bainier and its perfect stage *Byssosclamyces* Westling. *Transactions of the British Mycological Society*, 40(1), 17-IN13.
- Buenz, E., Bauer, B., Osmundson, T., & Motley, T. (2005). The traditional Chinese medicine *Cordyceps sinensis* and its effects on apoptotic homeostasis. *Journal of ethnopharmacology*, 96(1-2), 19-29.
- Bunyard, B. A., Nicholson, M. S., & Royse, D. J. (1994). A systematic assessment of *Morchella* using RFLP analysis of the 28S ribosomal RNA gene. *Mycologia*, 86(6), 762-772.
- Cesati, V. (1861). *Appunti per una futura Crittogamologia Insubrica*.
- Chen, Y.-Q., Hu, B., Xu, F., Zhang, W., Zhou, H., & Qu, L.-H. (2004). Genetic variation of *Cordyceps sinensis*, a fruit-body-producing entomopathogenic species from different geographical regions in China. *FEMS microbiology letters*, 230(1), 153-158.
- Chiu, C.-P., Liu, S.-C., Tang, C.-H., Chan, Y., El-Shazly, M., Lee, C.-L., . . . Wu, Y.-C. (2016). Anti-inflammatory cerebroside from cultivated *Cordyceps militaris*. *Journal of agricultural and food chemistry*, 64(7), 1540-1548.
- Choi, Y.-W., Hyde, K. D., & Ho, W. (1999). Single spore isolation of fungi. *Fungal diversity*.
- Da Silva, G. A., Lumini, E., Maia, L. C., Bonfante, P., & Bianciotto, V. (2006). Phylogenetic analysis of Glomeromycota by partial LSU rDNA sequences. *Mycorrhiza*, 16(3), 183-189.
- Dang, H.-N., Wang, C.-L., & Lay, H.-L. (2018). Effect of nutrition, vitamin, grains, and temperature on the mycelium growth and antioxidant capacity of *Cordyceps militaris* (strains AG-1 and PSJ-1). *Journal of radiation research and applied sciences*, 11(2), 130-138.

- Das, S. K., Masuda, M., Sakurai, A., & Sakakibara, M. (2010). Medicinal uses of the mushroom *Cordyceps militaris*: current state and prospects. *Fitoterapia*, 81(8), 961-968.
- De Bary, A. (1867). Zur kenntniss insectentodtender Pilze. *Bot Zeit*, 25, 1-7.
- Dong, J. Z., Lei, C., Ai, X. R., & Wang, Y. (2012). Selenium enrichment on *Cordyceps militaris* link and analysis on its main active components. *Applied biochemistry and biotechnology*, 166(5), 1215-1224.
- Dube, H. (2013). *An introduction to fungi*: Scientific Publishers.
- Eriksson, O., Baral, H., Currah, R., Hansen, K., Kurtzman, C., Læssøe, T., & Rambold, G. (2003). Notes on ascomycete systematics Nos 3580–3623. *Myconet*, 9, 91-103.
- Farlow, W. G., & Seymour, A. B. (1888). *A provisional host-index of the fungi of the United States* (Vol. 1).
- Gray, G. R. (1858). *Notices of insects that are known to form the bases of fungoid parasites*: Privately printed.
- Gu, H., & Liang, M. (1987). Study on the manual cultivation of *Cordyceps militaris*. *Med Inf Lett*, 5, 51-52.
- Guo, L.-X., Hong, Y.-H., Zhou, Q.-Z., Zhu, Q., Xu, X.-M., & Wang, J.-H. (2017). Fungus-larva relation in the formation of *Cordyceps sinensis* as revealed by stable carbon isotope analysis. *Scientific reports*, 7(1), 7789.
- Hao, D., Chen, S., Xiao, P., & Peng, Y. (2010). Authentication of medicinal plants by DNA-based markers and genomics. *Chin Herb Med*, 2(4), 250-261.
- Hitchcock, S. W. (1961). Pupal mortality of the orange-striped oakworm. *Journal of Economic Entomology*, 54(5), 962-964.
- Holliday, J. C., Cleaver, P., Loomis-Powers, M., & Patel, D. (2004). Analysis of quality and techniques for hybridization of medicinal fungus *Cordyceps sinensis* (Berk.) Sacc.(Ascomycetes). *International Journal of Medicinal Mushrooms*, 6(2), 151-164.
- Hong, I.-P., Kang, P.-D., Kim, K.-Y., Nam, S.-H., Lee, M.-Y., Choi, Y.-S., . . . Humber, R. A. (2010). Fruit body formation on silkworm by *Cordyceps militaris*. *Mycobiology*, 38(2), 128-132.
- Izgi, K., Iskender, B., Sakalar, C., Arslanhan, A., Saraymen, B., & Canatan, H. (2015). Evaluation of two different adjuvants with immunogenic uroplakin 3A-derived peptide for their ability to evoke an immune response in mice. *European cytokine network*, 26(2), 46-56.
- Jing, Y. (1987). A summary report on domestication of *Cordyceps militaris* strain. *Edible Fungi*, 2, 5-6.
- Joshi, S. P., Ranjekar, P. K., & Gupta, V. S. (1999). Molecular markers in plant genome analysis. *Curr Sci*, 77(2), 230-240.
- Kang, J., Liang, Z., Liu, A., & Kong, C. (2000). Molecular evidence of polymorphism in *Cordyceps* based on 5.8 S rDNA and ITS2 sequences. *Mycosystema*, 19(4), 492-497.
- Kang, P.-D., Sung, G.-B., Kim, K.-Y., Kim, M.-J., Hong, I.-P., & Ha, N.-G. (2010). Breeding of a silkworm variety for synnemata production of *Isaria tenuipes*. *Mycobiology*, 38(3), 180-183.

- Kim, S.-Y., Shrestha, B., Sung, G.-H., Han, S.-K., & Sung, J.-M. (2010). Optimum conditions for artificial fruiting body formation of *Cordyceps cardinalis*. *Mycobiology*, 38(2), 133-136.
- Kinjo, N., & Zang, M. (2001). Morphological and phylogenetic studies on *Cordyceps sinensis* distributed in southwestern China. *Mycoscience*, 42(6), 567-574.
- Kobayasi, Y. (1941). The genus *Cordyceps* and its allies. *Science Reports of the Tokyo Bunrika Daigaku*, 5, 53-260.
- Kobayasi, Y. (1982). Keys to the taxa of the genera *Cordyceps* and *Torrubiella*. *Transaction of the Mycological Society of Japan*.
- Kryukov, V. Y., Yaroslavtseva, O., Lednev, G., & Borisov, B. (2011). Local epizootics caused by teleomorphic cordycipitoid fungi (Ascomycota: Hypocreales) in populations of forest lepidopterans and sawflies of the summer-autumn complex in Siberia. *Microbiology*, 80(2), 286-295.
- Kuo, H.-C., Su, Y.-L., Yang, H.-L., & Chen, T.-Y. (2005). Identification of Chinese medicinal fungus *Cordyceps sinensis* by PCR-single-stranded conformation polymorphism and phylogenetic relationship. *Journal of agricultural and food chemistry*, 53(10), 3963-3968.
- Lagerberg, T. (1922). Smärre meddelanden: *Cordiceps militaris* (L.) Link i Sverige. *Svensk Bot Tidskr*, 16, 285-290.
- Lam, K., Chan, G., Xin, G.-Z., Xu, H., Ku, C.-F., Chen, J.-P., . . . Tsim, K. (2015). Authentication of *Cordyceps sinensis* by DNA analyses: Comparison of ITS sequence analysis and RAPD-derived molecular markers. *Molecules*, 20(12), 22454-22462.
- Leck, A. (1999). Preparation of lactophenol cotton blue slide mounts. *Community Eye Health*, 12(30), 24.
- Lee, B.-J., Lee, M., Kim, Y.-G., Lee, K.-W., Choi, Y.-S., Lee, B.-E., & Song, H.-Y. (2013). Cultural characteristics of *Cordyceps militaris* strain Yedang 3 on various media and nutritional conditions. *Journal of Mushroom*, 11(3), 124-130.
- Li, C., Li, Z., Fan, M., Cheng, W., Long, Y., Ding, T., & Ming, L. (2006). The composition of *Hirsutella sinensis*, anamorph of *Cordyceps sinensis*. *Journal of Food Composition and Analysis*, 19(8), 800-805.
- Li, Y., Jiao, L., & Yao, Y.-J. (2013). Non-concerted ITS evolution in fungi, as revealed from the important medicinal fungus *Ophiocordyceps sinensis*. *Molecular phylogenetics and evolution*, 68(2), 373-379.
- Liang, H.-H., Cheng, Z., Yang, X.-L., Li, S., Ding, Z.-Q., Zhou, T.-S., . . . Chen, J.-K. (2008). Genetic diversity and structure of *Cordyceps sinensis* populations from extensive geographical regions in China as revealed by inter-simple sequence repeat markers. *The Journal of Microbiology*, 46(5), 549-556.
- Lin, L.-T., Lai, Y.-J., Wu, S.-C., Hsu, W.-H., & Tai, C.-J. (2018). Optimal conditions for cordycepin production in surface liquid-cultured *Cordyceps militaris* treated with porcine liver extracts for suppression of oral cancer. *journal of food and drug analysis*, 26(1), 135-144.
- Liu, A. (1997). Determination of two species of *Cordyceps* by ascospore microcycle conidiation. *Mycosystema*, 16, 297-299.



- Liu, H.-j., Hu, H.-b., Chu, C., Li, Q., & Li, P. (2011). Morphological and microscopic identification studies of Cordyceps and its counterfeits. *Acta Pharmaceutica Sinica B*, 1(3), 189-195.
- Liu, J.-y., Feng, C.-p., Li, X., Chang, M.-c., Meng, J.-l., & Xu, L.-j. (2016). Immunomodulatory and antioxidative activity of Cordyceps militaris polysaccharides in mice. *International journal of biological macromolecules*, 86, 594-598.
- Liu, Z., Liang, Z., Whalley, A., Liu, A., & Yao, Y. (2001). A new species of Beauveria, the anamorph of Cordyceps sobolifera. *Fungal diversity*, 7, 61-70.
- Lord, N., Kaplan, C. W., Shank, P., Kitts, C. L., & Elrod, S. L. (2002). Assessment of fungal diversity using terminal restriction fragment (TRF) pattern analysis: comparison of 18S and ITS ribosomal regions. *FEMS Microbiology Ecology*, 42(3), 327-337.
- Ma, L., Zhang, S., & Du, M. (2015). Cordycepin from Cordyceps militaris prevents hyperglycemia in alloxan-induced diabetic mice. *Nutrition Research*, 35(5), 431-439.
- Ma, T., Feng, Y., Wu, X.-p., Zhang, Y., Ma, Y., & Wang, Z. (2007). Primary investigation of a host insect of Cordyceps militaris and analysis of its main ingredients. *FOREST RESEARCH-CHINESE ACADEMY OF FORESTRY*, 20(1), 63.
- Müller-Kögler, E. (1964). Cordyceps militaris (Fr.) Link: Beobachtungen und Versuche anlässlich eines Fundes auf Tipula paludosa Meig.(Dipt., Tipul.). *Zeitschrift für Angewandte Entomologie*, 55(1-4), 409-418.
- Panigrahi, A. (1995). Fungus Cordyceps militaris(Fries) link infestation in the pupa of the tea pest Andraca bipunctata Wlker. *Environment and ecology. Kalyani*, 13(4), 942-946.
- Park, J.-P., Kim, S. W., Hwang, H., & Yun, J.-W. (2001). Optimization of submerged culture conditions for the mycelial growth and exo-biopolymer production by Cordyceps militaris. *Letters in Applied Microbiology*, 33(1), 76-81.
- Park, Y.-J., & Min, B.-R. (2005). Sequence analysis of the internal transcribed spacer of ribosomal DNA in the genus rhizopus. *Mycobiology*, 33(2), 109-112.
- Pathania, P., Joshi, M., & Sagar, A. (2015). Morphological, Physiological and Molecular studies on wildlly collected Cordyceps militaris from North West Himalayas, India. *European Journal of Biotechnology and Bioscience*, 3(1), 53-62.
- Peay, K. G., Kennedy, P. G., & Bruns, T. D. (2008). Fungal community ecology: a hybrid beast with a molecular master. *Bioscience*, 58(9), 799-810.
- Petch, T. (1936). Cordyceps militaris and Isaria farinosa. *Transactions of the British Mycological Society*, 20(3-4), 216-224.
- Petch, T. (1948). A revised list of British entomogenous fungi. *Transactions of the British Mycological Society*, 31(3-4), 286-304.
- Pettit, R. H. (1895). *Studies in artificial cultures of entomogenous fungi* (Vol. 97): Cornell University.
- Qiwu, C. (1997). Study on Cordyceps (Fr.) fungi in Shennongjia forest district. *Hubei Agricultural Sciences*, 6.
- Raethong, N., Laoteng, K., & Vongsangnak, W. (2018). Uncovering global metabolic response to cordycepin production in Cordyceps militaris through



- transcriptome and genome-scale network-driven analysis. *Scientific reports*, 8(1), 9250.
- Redecker, D. (2002). Molecular identification and phylogeny of arbuscular mycorrhizal fungi *Diversity and Integration in Mycorrhizas* (pp. 67-73): Springer.
- Sangtian, T. (2009). Studies on mushrooms, fungi and molds in mangroves in Thailand *Bangkok, Office of the mangrove conservation*, 215.
- Sasaki, F., Miyamoto, T., Tamai, Y., & Yajima, T. (2007). Note on *Cordyceps brongniartii* Shimazu collected from the wild in Japan. *Mycoscience*, 48(5), 312-315.
- Sato, H., & Shimazu, M. (2002). Stromata production for *Cordyceps militaris* (Clavicipitales: Clavicipitaceae) by injection of hyphal bodies to alternative host insects. *Applied entomology and zoology*, 37(1), 85-92.
- Sato, H., Shimazu, M., & Kamata, N. (1994). Detection of *Cordyceps militaris* LINK (Clavicipitales: Clavicipitaceae) by Burying Pupae of *Quadricalcarifera punctatella* MOTSCHLSKY (Lepidoptera: Notodontidae). *Applied entomology and zoology*, 29(1), 130-132.
- Schoch, C. L., Seifert, K. A., Huhndorf, S., Robert, V., Spouge, J. L., Levesque, C. A., . . . Consortium, F. B. (2012). Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for Fungi. *Proceedings of the National Academy of Sciences*, 109(16), 6241-6246.
- Shrestha, B., Han, S.-K., Yoon, K.-S., & Sung, J.-M. (2005). Morphological characteristics of conidiogenesis in *Cordyceps militaris*. *Mycobiology*, 33(2), 69-76.
- Shrestha, B., Lee, W.-H., Han, S.-K., & Sung, J.-M. (2006). Observations on some of the mycelial growth and pigmentation characteristics of *Cordyceps militaris* isolates. *Mycobiology*, 34(2), 83-91.
- Shrestha, B., Zhang, W., Zhang, Y., & Liu, X. (2010). What is the Chinese caterpillar fungus *Ophiocordyceps sinensis* (Ophiocordycipitaceae)? *Mycology*, 1(4), 228-236.
- Simon, M., Binder, M., Adam, G., Hartig, A., & Ruis, H. (1992). Control of peroxisome proliferation in *Saccharomyces cerevisiae* by ADR1, SNF1 (CAT1, CCR1) and SNF4 (CAT3). *Yeast*, 8(4), 303-309.
- Sivichai, S. (2001). Capability of entomopathogenic isolates for the control of insect population. *Biosphere*, 3(3), 9-12.
- Song, J., Wang, Y., Teng, M., Zhang, S., Yin, M., Lu, J., . . . Teng, L. (2016). *Cordyceps militaris* induces tumor cell death via the caspase-dependent mitochondrial pathway in HepG2 and MCF-7 cells. *Molecular medicine reports*, 13(6), 5132-5140.
- Sopp, J. (1911). Untersuchungen über insektenvertilgende Pilze bei den letzten Kieferspinnerepidemien in Norwegen. *Vidensk. Selsk. Kria. Skr. I. Math.-naturv. Kl*, 2, 1-56.
- Strobel, G., Yang, X., Sears, J., Kramer, R., Sidhu, R. S., & Hess, W. (1996). Taxol from *Pestalotiopsis microspora*, an endophytic fungus of *Taxus wallachiana*. *Microbiology*, 142(2), 435-440.

- Sung, G.-H., Hywel-Jones, N. L., Sung, J.-M., Luangsa-ard, J. J., Shrestha, B., & Spatafora, J. W. (2007). Phylogenetic classification of Cordyceps and the clavicipitaceous fungi. *Studies in mycology*, 57, 5-59.
- Sung, G.-H., Shrestha, B., Han, S.-K., Kim, S.-Y., & Sung, J.-M. (2010). Growth and cultural characteristics of Cordyceps cardinalis collected from Korea. *Mycobiology*, 38(4), 274-281.
- Sung, G.-H., Shrestha, B., Han, S.-K., & Sung, J.-M. (2011). Growth and cultural characteristics of Ophiocordyceps longissima collected in Korea. *Mycobiology*, 39(2), 85-91.
- Sung, G.-H., Spatafora, J. W., Zare, R., Hodge, K. T., & Gams, W. (2001). A revision of Verticillium sect. Prostrata. II. Phylogenetic analyses of SSU and LSU nuclear rDNA sequences from anamorphs and teleomorphs of the Clavicipitaceae. *Nova Hedwigia*, 72(3-4), 311-328.
- Sung, J.-M., Choi, Y.-S., Shrestha, B., & Park, Y.-J. (2002). Investigation on artificial fruiting of Cordyceps militaris. *The Korean Journal of Mycology*, 30(1), 6-10.
- Sung, J.-M., Park, Y.-J., Lee, J.-O., Han, S.-K., Lee, W.-H., Choi, S.-K., & Shrestha, B. (2006). Selection of superior strains of Cordyceps militaris with enhanced fruiting body productivity. *Mycobiology*, 34(3), 131-137.
- Suwanpinit, N., & Suwanpinit, P. (1998). General microbiology Bangkok, Chulalongkorn university.
- Sylvia S. Mader, W., Hocker, J., Arthur, J., Petty, J. W., Martocchio, J. J., Cheeseman, H. R., . . . Rosania, R. J. (2007). *Interpersonal Conflict 9th Edition*: McGraw-Hill Higher Education.
- Tapingkae, T. (2012). Cordyceps mushroom cultivation Bangkok, two-four printing, 15-20.
- Tapingkae, T. (2013). Analysis of the Cordycepin in Cordyceps militaris cultivation by rice ChiangMai, ChiangMai Rajabhat University.
- Tapingkae, T. (2014). Study on Cultivation and Utilization of Medicinal Mushroom Cordyceps militaris ChiangMai, ChiangMai Rajabhat University.
- Tatani, K., Hiratochi, M., Kikuchi, N., Kuramochi, Y., Watanabe, S., Yamauchi, Y., . . . Shuto, S. (2016). Identification of adenine and benzimidazole nucleosides as potent human concentrative nucleoside transporter 2 inhibitors: potential treatment for hyperuricemia and gout. *Journal of medicinal chemistry*, 59(8), 3719-3731.
- Tianzhu, Z., Shihai, Y., & Juan, D. (2015). The effects of cordycepin on ovalbumin-induced allergic inflammation by strengthening Treg response and suppressing Th17 responses in ovalbumin-sensitized mice. *Inflammation*, 38(3), 1036-1043.
- Tulasne, L. R. T., C. (1865). Selecta Fungorum Carpologia: Nectriei- Phacidiei- Pezizei. Paris, 3.
- Ulvinen, T. (1969). Hyönteisissä elävistä Cordyceps-lajeista eräistä kuroma-asteisista sienistä. Pikkutietoja-Notulae. Paper presented at the Annales Entomologici Fennici.
- Wang, L., Zhang, W., Hu, B., Chen, Y., & Qu, L. (2008). Genetic variation of Cordyceps militaris and its allies based on phylogenetic analysis of rDNA ITS sequence data. *Fungal Divers*, 31, 147-155.

- Watanabe, M., Yonezawa, T., Lee, K.-i., Kumagai, S., Sugita-Konishi, Y., Goto, K., & Hara-Kudo, Y. (2011). Molecular phylogeny of the higher and lower taxonomy of the *Fusarium* genus and differences in the evolutionary histories of multiple genes. *BMC Evolutionary Biology*, 11(1), 322.
- Wen, T.-C., Zhu, R.-C., Kang, J.-C., Huang, M.-H., Tan, D.-B., Ariyawansa, H., . . . Liu, H. (2013). *Ophiocordyceps xuefengensis* sp. nov. from larvae of *Phassus nodus* (Hepialidae) in Hunan Province, southern China. *Phytotaxa*, 123(1), 41-50.
- Wen, T., Li, G., Kang, J., Kang, C., & Hyde, K. D. (2014). Optimization of solid-state fermentation for fruiting body growth and cordycepin production by *Cordyceps militaris*. *Chiang Mai J Sci*, 41(4), 858-872.
- Wen, T., Long, F., Kang, C., Wang, F., & Zeng, W. (2017). Effects of additives and bioreactors on cordycepin production from *Cordyceps militaris* in liquid static culture. *Mycosphere*, 8(7), 886-898.
- White, T. J., Bruns, T., Lee, S., & Taylor, J. (1990). Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. *PCR protocols: a guide to methods and applications*, 18(1), 315-322.
- Xiong, C., Xia, Y., Zheng, P., Shi, S., & Wang, C. (2010). Developmental stage-specific gene expression profiling for a medicinal fungus *Cordyceps militaris*. *Mycology*, 1(1), 25-66.
- Yong, T., Zhang, M., Chen, D., Shuai, O., Chen, S., Su, J., . . . Xie, Y. (2016). Actions of water extract from *Cordyceps militaris* in hyperuricemic mice induced by potassium oxonate combined with hypoxanthine. *Journal of ethnopharmacology*, 194, 403-411.
- Yue-Qin, C., Ning, W., Hui, Z., & Liang-Hu, Q. (2002). Differentiation of medicinal *Cordyceps* species by rDNA ITS sequence analysis. *Planta medica*, 68(07), 635-639.
- Zhang, G., & Liang, Y. (2013). Improvement of fruiting body production in *Cordyceps militaris* by molecular assessment. *Archives of microbiology*, 195(8), 579-585.
- Zhang, P., Huang, C., Fu, C., Tian, Y., Hu, Y., Wang, B., . . . Song, E. (2015). Cordycepin (3'-deoxyadenosine) suppressed HMGA2, Twist1 and ZEB1-dependent melanoma invasion and metastasis by targeting miR-33b. *Oncotarget*, 6(12), 9834.
- Zhong, X., Peng, Q., Qi, L., Lei, W., & Liu, X. (2010). rDNA-targeted PCR primers and FISH probe in the detection of *Ophiocordyceps sinensis* hyphae and conidia. *Journal of microbiological methods*, 83(2), 188-193.

## APPENDIX

**Appendix 1** Groups aligned by sequence fragments (approximately 548 and 1223) generated from ITS and 28S gene respectively, amplified by using DNA template from seven strains *C. militaris*

### ITS gene

#### >Cmi1

AAGTTCAGCGGGTATTCCTACCTGATCCGAGGTCAACGTTTCAGAGTTGGG  
CGTTTTACGGCGTGGCCACGTCGGGTTCCTCGGTGCGAGTTGGAGTACTAC  
GCAGAGGTCGCCGCGGACGGGCGCCACTTCATTTCTGGGGGCGGCGGTGT  
GCTGCCGGTCCCCAACGCCGACATCCCCAGGGGACGTCGAGGGTTGAAA  
TGACGCTCGAACAGGCATGCCCCGCCAGAATGCTGGCGGGCGCAATGTGCG  
TTCAAAGATTCGATGATTCACTGAATTCTGCAATTCACATTACTTATCGCA  
TTTCGCTGCGTTCTTCATCGATGCCAGAGCCAAGAGATCCGTTGTTGAAAG  
TTTTGATTCATTTGTTTTGCCTTGCGGCGGATTCAGAAAACTGGTAGATA  
CAGTGTTTGGGGCCCCCGACGGCCGCCGCCAGGCCCCGCGTCCAGGCGCT  
GGGCGAGTCCGCCGAAGCAACGATAGGTATGTTTACAAAGGGTTGGGAGT  
TGAAAACTCGTTAATGATCCCTCCGCTGGTTCACCAACGGAGAC

#### >Cmi2

AAGTTCAGCGGGTATTCCTACCTGATCCGAGGTCAACGTTTCAGAGTTGGG  
CGTTTTACGGCGTGGCCACGTCGGGTTCCTCGGTGCGAGTTGGAGTACTAC  
GCAGAGGTCGCCGCGGACGGGCGCCACTTCATTTCTGGGGGCGGCGGTGT  
GCTGCCGGTCCCCAACGCCGACATCCCCAGGGGACGTCGAGGGTTGAAA  
TGACGCTCGAACAGGCATGCCCCGCCAGAATGCTGGCGGGCGCAATGTGCG  
TTCAAAGATTCGATGATTCACTGAATTCTGCAATTCACATTACTTATCGCA  
TTTCGCTGCGTTCTTCATCGATGCCAGAGCCAAGAGATCCGTTGTTGAAAG  
TTTTGATTCATTTGTTTTGCCTTGCGGCGGATTCAGAAAACTGGTAGATA  
CAGTGTTTGGGGCCCCCGACGGCCGCCGCCAGGCCCCGCGTCCAGGCGCT  
GGGCGAGTCCGCCGAAGCAACGATAGGTATGTTTACAAAGGGTTGGGAGT  
TGAAAACTCGTTAATGATCCCTCCGCTGGTTCACCAACGGAGAC

**>Cmi3**

AAGTTCAGCGGGTATTCCTACCTGATCCGAGGTCAACGTTTCAGAGTTGGG  
CGTTTTACGGCGTGGCCACGTCGGGTTCCTGGTGCGAGTTGGAGTACTAC  
GCAGAGGTCGCCGCGGACGGGCGCCACTTCATTTCTGGGGGCGGCGGTGT  
GCTGCCGGTCCCCAACGCCGACATCCCCCAGGGGACGTCGAGGGTTGAAA  
TGACGCTCGAACAGGCATGCCCCGCCAGAATGCTGGCGGGCGCAATGTGCG  
TTCAAAGATTCGATGATTCACTGAATTCTGCAATTCACATTACTTATCGCA  
TTTCGCTGCGTTCTTCATCGATGCCAGAGCCAAGAGATCCGTTGTTGAAAG  
TTTTGATTCATTTGTTTTGCCTTGCGGCGGATTCAGAAAACTGGTAGATA  
CAGTGTTTGGGGCCCCCGACGGCCGCGGCCAGGCCCCGCGTCCAGGCGCT  
GGGCGAGTCCGCCGAAGCAACGATAGGTATGTTTACAAAGGGTTGGGAGT  
TGAAAACTCGTTAATGATCCCTCCGCTGGTTCACCAACGGAGAC

**>Cmi4**

AAGTTCAGCGGGTATTCCTACCTGATCCGAGGTCAACGTTTCAGAGTTGGG  
CGTTTTACGGCGTGGCCACGTCGGGTTCCTGGTGCGAGTTGGAGTACTAC  
GCAGAGGTCGCCGCGGACGGGCGCCACTTCATTTCTGGGGGCGGCGGTGT  
GCTGCCGGTCCCCAACGCCGACATCCCCCAGGGGACGTCGAGGGTTGAAA  
TGACGCTCGAACAGGCATGCCCCGCCAGAATGCTGGCGGGCGCAATGTGCG  
TTCAAAGATTCGATGATTCACTGAATTCTGCAATTCACATTACTTATCGCA  
TTTCGCTGCGTTCTTCATCGATGCCAGAGCCAAGAGATCCGTTGTTGAAAG  
TTTTGATTCATTTGTTTTGCCTTGCGGCGGATTCAGAAAACTGGTAGATA  
CAGTGTTTGGGGCCCCCGACGGCCGCGGCCAGGCCCCGCGTCCAGGCGCT  
GGGCGAGTCCGCCGAAGCAACGATAGGTATGTTTACAAAGGGTTGGGAGT  
TGAAAACTCGTTAATGATCCCTCCGCTGGTTCACCAACGGAGAC



**>Cmi5**

AAGTTCAGCGGGTATTCCTACCTGATCCGAGGTCAACGTTTCAGAGTTGGG  
CGTTTTACGGCGTGGCCACGTCGGGTTCCTCGGTGCGAGTTGGAGTACTAC  
GCAGAGGTCGCCGCGGACGGGCGCCACTTCATTTTCGGGGGCGGCGGTGT  
GCTGCCGGTCCCCAACGCCGACATCCCCAGGGGACGTCGAGGGTTGAAA  
TGACGCTCGAACAGGCATGCCCCGCCAGAATGCTGGCGGGCGCAATGTGCG  
TTCAAAGATTCGATGATTCACCTGAATTCTGCAATTCACATTACTTATCGCA  
TTTCGCTGCGTTCTTCATCGATGCCAGAGCCAAGAGATCCGTTGTTGAAAG  
TTTTGATTCATTTGTTTTGCCTTGCGGCGGATTCAGAAAACTGGTAGATA  
CAGTGTTTGGGGCCCCCGACGGCCGCGGCCAGGCCCCGCGTCCAGGCGCT  
GGGCGAGTCCGCCGAAGCAACGATAGGTATGTTTACAAAGGGTTGGGAGT  
TGAAAACTCGTTAATGATCCCTCCGCTGGTTCACCAACGGAGAC

**>Cmi6**

AAGTTCAGCGGGTATTCCTACCTGATCCGAGGTCAACGTTTCAGAGTTGGG  
CGTTTTACGGCGTGGCCACGTCGGGTTCCTCGGTGCGAGTTGGAGTACTAC  
GCAGAGGTCGCCGCGGACGGGCGCCACTTCATTTTCGGGGGCGGCGGTGT  
GCTGCCGGTCCCCAACGCCGACATCCCCAGGGGACGTCGAGGGTTGAAA  
TGACGCTCGAACAGGCATGCCCCGCCAGAATGCTGGCGGGCGCAATGTGCG  
TTCAAAGATTCGATGATTCACCTGAATTCTGCAATTCACATTACTTATCGCA  
TTTCGCTGCGTTCTTCATCGATGCCAGAGCCAAGAGATCCGTTGTTGAAAG  
TTTTGATTCATTTGTTTTGCCTTGCGGCGGATTCAGAAAACTGGTAGATA  
CAGTGTTTGGGGCCCCCGACGGCCGCGGCCAGGCCCCGCGTCCAGGCGCT  
GGGCGAGTCCGCCGAAGCAACGATAGGTATGTTTACAAAGGGTTGGGAGT  
TGAAAACTCGTTAATGATCCCTCCGCTGGTTCACCAACGGAGAC



**>Cmi7**

AAGTTCAGCGGGTATTCCTACCTGATCCGAGGTCAACGTTTCAGAGTTGGG  
CGTTTTACGGCGTGGCCACGTCGGGTCCCCGGTGCGAGTTGGAGTACTAC  
GCAGAGGTTCGCCGCGGACGGGGCCGCCACTTCATTTTCGGGGGCGGCGGTGT  
GCTGCCGGTCCCCAACGCCGACATCCCCCAGGGGACGTCGAGGGTTGAAA  
TGACGCTCGAACAGGCATGCCCCGCCAGAATGCTGGCGGGCGCAATGTGCG  
TTCAAAGATTCGATGATTCACTGAATTCTGCAATTCACATTACTTATCGCA  
TTTCGCTGCGTTCTTCATCGATGCCAGAGCCAAGAGATCCGTTGTTGAAAG  
TTTTGATTCATTTGTTTTGCCTTGCGGCGGATTCAGAAAACTGGTAGATA  
CAGTGTTTGGGGCCCCCGACGGCCGCCGCCAGGCCCCGCGTCCAGGCGCT  
GGGCGAGTCCGCCGAAGCAACGATAGGTATGTTTACAAAGGGTTGGGAGT  
TGAAAACTCGTTAATGATCCCTCCGCTGGTTCACCAACGGAGAC



**28S gene****>Cmi1**

GAAATCTGGCCCCCGGGTCCGAGTTGTAATTTGTAGAGGATGCTTTTGGCG  
AGGTGCCTTCCGAGTTCCCTGGAACGGGACGCCATAGAGGGTGAGAGCCC  
CGTCTGGTCGGACACCGAGCCTCTGTAAAGCTCCTTCGACGAGTCGAGTA  
GTTTGGGAATGCTGCTCAAAATGGGAGGTATATGTCTTCTAAAGCTAAAT  
ATTGGCCAGAGACCGATAGCGCACAAAGTAGAGTGATCGAAAGATGAAAA  
GCACTTTGAAAAGAGGGTTAAAAAGTACGTGAAATTGTTGAAAGGGAAG  
CGCCTATGACCAGACTTGGGCCCCGGTGAATCACCCAGCGTTCTCGCTGGT  
GCACTTCGCCGGGCACAGGCCAGCATCAGTTTGGCGCGGGGGAGAAAGG  
CTTCGGGAATGTGGCTCCCCTGGGAGTGTTATAGCCCGTTGCGCAATACCC  
TGCGCTGGACTGAGGTACGCGCATTCGCAAGGATGCTGGCGTAATGGTCA  
TCAGCGACCCGTCTTGAAACACGGACCAAGGAGTCGTCTTCGTATGCGAG  
TGTTTCGGGTGTCAAACCCCTACGCGTAATGAAAGTGAACGCAGGTGAGAG  
CTTCGGCGCATCATCGACCGATCCTGATGTTCTCGGATGGATTTGAGTAAG  
AGCATAACGGGGCCGGACCCGAAAGAAGGTGAACTATGCCTGTATAGGGT  
GAAGCCAGAGGAAACTCTGGTGGAGGCTCGCAGCGGTTCTGACGTGCGAA  
TCGATCGTCAAATATGGGCATGGGGGCGAAAGACTAATCGAACCTTCTAG  
TAGCTGGTTTCCGCCGAAGTTTCCCTCAGGATAGCAGTGTTGAACTCAGTT  
TTATGAGGTAAAGCGAATGATTAGGGACTCGGGGGCGCTATATTGCCTTC  
ATCCATTCTCAAACCTTTAAATATGTAAGAAGCCCTTGTTACTTGATTGAAC  
GTGGGCATTTCGAATGTATCAACACTAGTGGGCCATTTTTGGTAAGCAGAA  
CTGGCGATGCGGGATGAACCGAACGCGAGGTTAAGGTGCCGGAGTGAC  
GCTCATCAGACACCACAAAAGGTGTTAGTACATCTTGACAGCAGGACGGT  
GGCCATGGAAGTCGGAATCCGCTAAGGACTGTGTAACAACTCACCTGCCG  
AATGTACTAGCCCTGAAAATGGATGGCGCTCAAGCGTCCCACCCATACCT  
CGCCCTCAGGGTAGAAACGAAGC

**>Cmi2**

GAAATCTGGCCCCCGGGTCCGAGTTGTAATTTGTAGAGGATGCTTTTGGCG  
AGGTGCCTTCCGAGTTCCTTGGAACGGGACGCCATAGAGGGTGAGAGCCC  
CGTCTGGTTCGGACACCGAGCCTCTGTAAAGCTCCTTCGACGAGTCGAGTA  
GTTTGGGAATGCTGCTCAAAATGGGAGGTATATGTCTTCTAAAGCTAAAT  
ATTGGCCAGAGACCGATAGCGCACAAAGTAGAGTGATCGAAAGATGAAAA  
GCACTTTGAAAAGAGGGTTAAAAAGTACGTGAAATTGTTGAAAGGGAAG  
CGCCTATGACCAGACTTGGGCCCGGTGAATCACCCAGCGTTCTCGCTGGT  
GCACTTCGCCGGGCACAGGCCAGCATCAGTTTGGCGCGGGGGAGAAAGG  
CTTCGGGAATGTGGCTCCCCTGGGAGTGTTATAGCCCGTTGCGCAATACCC  
TGCGCTGGACTGAGGTACGCGCATTCGCAAGGATGCTGGCGTAATGGTCA  
TCAGCGACCCGTCTTGAAACACGGACCAAGGAGTCGTCTTCGTATGCGAG  
TGTTTCGGGTGTCAAACCCCTACGCGTAATGAAAGTGAACGCAGGTGAGAG  
CTTCGGCGCATCATCGACCGATCCTGATGTTCTCGGATGGATTTGAGTAAG  
AGCATACGGGGCCGGACCCGAAAGAAGGTGAACTATGCCTGTATAGGGT  
GAAGCCAGAGGAAACTCTGGTGGAGGCTCGCAGCGGTTCTGACGTGCGAA  
TCGATCGTCAAATATGGGCATGGGGGCGAAAGACTAATCGAACCTTCTAG  
TAGCTGGTTTCCGCCGAAGTTTCCCTCAGGATAGCAGTGTTGAACTCAGTT  
TTATGAGGTAAAGCGAATGATTAGGGACTCGGGGGCGCTATATTGCCTTC  
ATCCATTCTCAAACCTTTAAATATGTAAGAAGCCCTTGTTACTTGATTGAAC  
GTGGGCATTTCGAATGTATCAACACTAGTGGGCCATTTTTGGTAAGCAGAA  
CTGGCGATGCGGGATGAACCGAACGCGAGGTTAAGGTGCCGGAGTGGAC  
GCTCATCAGACACCACAAAAGGTGTTAGTACATCTTGACAGCAGGACGGT  
GGCCATGGAAGTCGGAATCCGCTAAGGACTGTGTAACTCACCTGCCG  
AATGTACTAGCCCTGAAAATGGATGGCGCTCAAGCGTCCCACCCATACCT  
CGCCCTCAGGGTAGAAACGAAGC

**>Cmi3**

GAAATCTGGCCCCCGGGTCCGAGTTGTAATTTGTAGAGGATGCTTTTGGCG  
AGGTGCCTTCCGAGTTCCTTGGAACGGGACGCCATAGAGGGTGAGAGCCC  
CGTCTGGTTCGGACACCGAGCCTCTGTAAAGCTCCTTCGACGAGTCGAGTA  
GTTTGGGAATGCTGCTCAAAATGGGAGGTATATGTCTTCTAAAGCTAAAT  
ATTGGCCAGAGACCGATAGCGCACAAAGTAGAGTGATCGAAAGATGAAAA  
GCACTTTGAAAAGAGGGTTAAAAAGTACGTGAAATTGTTGAAAGGGAAG  
CGCCTATGACCAGACTTGGGCCCGGTGAATCACCCAGCGTTCTCGCTGGT  
GCACTTCGCCGGGCACAGGCCAGCATCAGTTTGGCGCGGGGGAGAAAGG  
CTTCGGGAATGTGGCTCCCCTGGGAGTGTTATAGCCCGTTGCGCAATACCC  
TGCGCTGGACTGAGGTACGCGCATTCGCAAGGATGCTGGCGTAATGGTCA  
TCAGCGACCCGTCTTGAAACACGGACCAAGGAGTCGTCTTCGTATGCGAG  
TGTTTCGGGTGTCAAACCCCTACGCGTAATGAAAGTGAACGCAGGTGAGAG  
CTTCGGCGCATCATCGACCGATCCTGATGTTCTCGGATGGATTTGAGTAAG  
AGCATACGGGGCCGGACCCGAAAGAAGGTGAACTATGCCTGTATAGGGT  
GAAGCCAGAGGAAACTCTGGTGGAGGCTCGCAGCGGTTCTGACGTGCGAA  
TCGATCGTCAAATATGGGCATGGGGGCGAAAGACTAATCGAACCTTCTAG  
TAGCTGGTTTCCGCCGAAGTTTCCCTCAGGATAGCAGTGTTGAACTCAGTT  
TTATGAGGTAAAGCGAATGATTAGGGACTCGGGGGCGCTATATTGCCTTC  
ATCCATTCTCAAACCTTTAAATATGTAAGAAGCCCTTGTTACTTGATTGAAC  
GTGGGCATTTCGAATGTATCAACACTAGTGGGCCATTTTTGGTAAGCAGAA  
CTGGCGATGCGGGATGAACCGAACGCGAGGTTAAGGTGCCGGAGTGGAC  
GCTCATCAGACACCACAAAAGGTGTTAGTACATCTTGACAGCAGGACGGT  
GGCCATGGAAGTCGGAATCCGCTAAGGACTGTGTAACAACTCACCTGCCG  
AATGTACTAGCCCTGAAAATGGATGGCGCTCAAGCGTCCCACCCATACCT  
CGCCCTCAGGGTAGAAACGAAGC

**>Cmi4**

GAAATCTGGCCCCCGGGTCCGAGTTGTAATTTGTAGAGGATGCTTTTGGCG  
AGGTGCCTTCCGAGTTCCTTGGAACGGGACGCCATAGAGGGTGAGAGCCC  
CGTCTGGTTCGGACACCGAGCCTCTGTAAAGCTCCTTCGACGAGTCGAGTA  
GTTTGGGAATGCTGCTCAAAATGGGAGGTATATGTCTTCTAAAGCTAAAT  
ATTGGCCAGAGACCGATAGCGCACAAAGTAGAGTGATCGAAAGATGAAAA  
GCACTTTGAAAAGAGGGTTAAAAAGTACGTGAAATTGTTGAAAGGGAAG  
CGCCTATGACCAGACTTGGGCCCGGTGAATCACCCAGCGTTCTCGCTGGT  
GCACTTCGCCGGGCACAGGCCAGCATCAGTTTGGCGCGGGGGAGAAAGG  
CTTCGGGAATGTGGCTCCCCTGGGAGTGTTATAGCCCGTTGCGCAATACCC  
TGCGCTGGACTGAGGTACGCGCATTCGCAAGGATGCTGGCGTAATGGTCA  
TCAGCGACCCGTCTTGAAACACGGACCAAGGAGTCGTCTTCGTATGCGAG  
TGTTTCGGGTGTCAAACCCCTACGCGTAATGAAAGTGAACGCAGGTGAGAG  
CTTCGGCGCATCATCGACCGATCCTGATGTTCTCGGATGGATTTGAGTAAG  
AGCATACGGGGCCGGACCCGAAAGAAGGTGAACTATGCCTGTATAGGGT  
GAAGCCAGAGGAAACTCTGGTGGAGGCTCGCAGCGGTTCTGACGTGCGAA  
TCGATCGTCAAATATGGGCATGGGGGCGAAAGACTAATCGAACCTTCTAG  
TAGCTGGTTTCCGCCGAAGTTTCCCTCAGGATAGCAGTGTTGAACTCAGTT  
TTATGAGGTAAAGCGAATGATTAGGGACTCGGGGGCGCTATATTGCCTTC  
ATCCATTCTCAAACCTTTAAATATGTAAGAAGCCCTTGTTACTTGATTGAAC  
GTGGGCATTCTGAATGTATCAACACTAGTGGGCCATTTTTGGTAAGCAGAA  
CTGGCGATGCGGGATGAACCGAACGCGAGGTTAAGGTGCCGGAGTGGAC  
GCTCATCAGACACCACAAAAGGTGTTAGTACATCTTGACAGCAGGACGGT  
GGCCATGGAAGTCGGAATCCGCTAAGGACTGTGTAACTCACCTGCCG  
AATGTACTAGCCCTGAAAATGGATGGCGCTCAAGCGTCCCACCCATACCT  
CGCCCTCAGGGTAGAAACGAAGC



**>Cmi5**

GAAATCTGGCCCCCGGGTCCGAGTTGTAATTTGTAGAGGATGCTTTTGGCG  
AGGTGCCTTCCGAGTTCCTGGAACGGGACGCCATAGAGGGTGAGAGCCC  
CGTCTGGTTCGGACACCGAGCCTCTGTAAAGCTCCTTCGACGAGTCGAGTA  
GTTTGGGAATGCTGCTCAAAATGGGAGGTATATGTCTTCTAAAGCTAAAT  
ATTGGCCAGAGACCGATAGCGCACAAAGTAGAGTGATCGAAAGATGAAAA  
GCACTTTGAAAAGAGGGTTAAAAAGTACGTGAAATTGTTGAAAGGGAAG  
CGCCTATGACCAGACTTGGGCCCGGTGAATCACCCAGCGTTCTCGCTGGT  
GCACTTCGCCGGGCACAGGCCAGCATCAGTTTGGCGCGGGGGAGAAAGG  
CTTCGGGAATGTGGCTCCCCTGGGAGTGTTATAGCCCGTTGCGCAATACCC  
TGCGCTGGACTGAGGTACGCGCATTCGCAAGGATGCTGGCGTAATGGTCA  
TCAGCGACCCGTCTTGAAACACGGACCAAGGAGTCGTCTTCGTATGCGAG  
TGTTTCGGGTGTCAAACCCCTACGCGTAATGAAAGTGAACGCAGGTGAGAG  
CTTCGGCGCATCATCGACCGATCCTGATGTTCTCGGATGGATTTGAGTAAG  
AGCATACGGGGCCGGACCCGAAAGAAGGTGAACTATGCCTGTATAGGGT  
GAAGCCAGAGGAAACTCTGGTGGAGGCTCGCAGCGGTTCTGACGTGCGAA  
TCGATCGTCAAATATGGGCATGGGGGCGAAAGACTAATCGAACCTTCTAG  
TAGCTGGTTTCCGCCGAAGTTTCCCTCAGGATAGCAGTGTTGAACTCAGTT  
TTATGAGGTAAAGCGAATGATTAGGGACTCGGGGGCGCTATATTGCCTTC  
ATCCATTCTCAAACCTTTAAATATGTAAGAAGCCCTTGTTACTTGATTGAAC  
GTGGGCATTTCGAATGTATCAACACTAGTGGGCCATTTTTGGTAAGCAGAA  
CTGGCGATGCGGGATGAACCGAACGCGAGGTTAAGGTGCCGGAGTGGAC  
GCTCATCAGACACCACAAAAGGTGTTAGTACATCTTGACAGCAGGACGGT  
GGCCATGGAAGTCGGAATCCGCTAAGGACTGTGTAACAACTCACCTGCCG  
AATGTACTAGCCCTGAAAATGGATGGCGCTCAAGCGTCCCACCCATACCT  
CGCCCTCAGGGTAGAAACGAAGC

**>Cmi6**

GAAATCTGGCCCCCGGGTCCGAGTTGTAATTTGTAGAGGATGCTTTTGGCG  
AGGTGCCTTCCGAGTTCCTTGGAACGGGACGCCATAGAGGGTGAGAGCCC  
CGTCTGGTTCGGACACCGAGCCTCTGTAAAGCTCCTTCGACGAGTCGAGTA  
GTTTGGGAATGCTGCTCAAAATGGGAGGTATATGTCTTCTAAAGCTAAAT  
ATTGGCCAGAGACCGATAGCGCACAAAGTAGAGTGATCGAAAGATGAAAA  
GCACTTTGAAAAGAGGGTTAAAAAGTACGTGAAATTGTTGAAAGGGAAG  
CGCCTATGACCAGACTTGGGCCCGGTGAATCACCCAGCGTTCTCGCTGGT  
GCACTTCGCCGGGCACAGGCCAGCATCAGTTTGGCGCGGGGGAGAAAGG  
CTTCGGGAATGTGGCTCCCCTGGGAGTGTTATAGCCCGTTGCGCAATACCC  
TGCGCTGGACTGAGGTACGCGCATTCGCAAGGATGCTGGCGTAATGGTCA  
TCAGCGACCCGTCTTGAAACACGGACCAAGGAGTCGTCTTCGTATGCGAG  
TGTTTCGGGTGTCAAACCCCTACGCGTAATGAAAGTGAACGCAGGTGAGAG  
CTTCGGCGCATCATCGACCGATCCTGATGTTCTCGGATGGATTTGAGTAAG  
AGCATACGGGGCCGGACCCGAAAGAAGGTGAACTATGCCTGTATAGGGT  
GAAGCCAGAGGAAACTCTGGTGGAGGCTCGCAGCGGTTCTGACGTGCGAA  
TCGATCGTCAAATATGGGCATGGGGGCGAAAGACTAATCGAACCTTCTAG  
TAGCTGGTTTCCGCCGAAGTTTCCCTCAGGATAGCAGTGTTGAACTCAGTT  
TTATGAGGTAAAGCGAATGATTAGGGACTCGGGGGCGCTATATTGCCTTC  
ATCCATTCTCAAACCTTTAAATATGTAAGAAGCCCTTGTTACTTGATTGAAC  
GTGGGCATTCTGAATGTATCAACACTAGTGGGCCATTTTTGGTAAGCAGAA  
CTGGCGATGCGGGATGAACCGAACGCGAGGTTAAGGTGCCGGAGTGGAC  
GCTCATCAGACACCACAAAAGGTGTTAGTACATCTTGACAGCAGGACGGT  
GGCCATGGAAGTCGGAATCCGCTAAGGACTGTGTAACAACTCACCTGCCG  
AATGTACTAGCCCTGAAAATGGATGGCGCTCAAGCGTCCCACCCATACCT  
CGCCCTCAGGGTAGAAACGAAGC

**>Cmi7**

GAAATCTGGCCCCCGGGTCCGAGTTGTAATTTGTAGAGGATGCTTTTGGCG  
AGGTGCCTTCCGAGTTCCTGGAACGGGACGCCATAGAGGGTGAGAGCCC  
CGTCTGGTTCGGACACCGAGCCTCTGTAAAGCTCCTTCGACGAGTCGAGTA  
GTTTGGGAATGCTGCTCAAAATGGGAGGTATATGTCTTCTAAAGCTAAAT  
ATTGGCCAGAGACCGATAGCGCACAAGTAGAGTGATCGAAAGATGAAAA  
GCACTTTGAAAAGAGGGTTAAAAAGTACGTGAAATTGTTGAAAGGGAAG  
CGCCTATGACCAGACTTGGGCCCGGTGAATCACCCAGCGTTCTCGCTGGT  
GCACTTCGCCGGGCACAGGCCAGCATCAGTTTGGCGCGGGGGAGAAAGG  
CTTCGGGAATGTGGCTCCCCTGGGAGTGTTATAGCCCGTTGCGCAATACCC  
TGCGCTGGACTGAGGTACGCGCATTCGCAAGGATGCTGGCGTAATGGTCA  
TCAGCGACCCGTCTTGAAACACGGACCAAGGAGTCGTCTTCGTATGCGAG  
TGTTTCGGGTGTCAAACCCCTACGCGTAATGAAAGTGAACGCAGGTGAGAG  
CTTCGGCGCATCATCGACCGATCCTGATGTTCTCGGATGGATTTGAGTAAG  
AGCATACGGGGCCGGACCCGAAAGAAGGTGAACTATGCCTGTATAGGGT  
GAAGCCAGAGGAAACTCTGGTGGAGGCTCGCAGCGGTTCTGACGTGCGAA  
TCGATCGTCAAATATGGGCATGGGGGCGAAAGACTAATCGAACCTTCTAG  
TAGCTGGTTTCCGCCGAAGTTTCCCTCAGGATAGCAGTGTTGAACTCAGTT  
TTATGAGGTAAAGCGAATGATTAGGGACTCGGGGGCGCTATATTGCCTTC  
ATCCATTCTCAAACCTTTAAATATGTAAGAAGCCCTTGTTACTTGATTGAAC  
GTGGGCATTCTGAATGTATCAACACTAGTGGGCCATTTTTGGTAAGCAGAA  
CTGGCGATGCGGGATGAACCGAACGCGAGGTTAAGGTGCCGGAGTGGAC  
GCTCATCAGACACCACAAAAGGTGTTAGTACATCTTGACAGCAGGACGGT  
GGCCATGGAAGTCGGAATCCGCTAAGGACTGTGTAACAACTCACCTGCCG  
AATGTACTAGCCCTGAAAATGGATGGCGCTCAAGCGTCCCACCCATACCT  
CGCCCTCAGGGTAGAAACGAAGC